

Practice Problems 12 Monday, November 25, 2024

1. Let D_n be the determinant of order n of which the element in the *i*th row and the *j*th column is the absolute value of the difference of i and j. Show that D_n is equal to

$$(-1)^{n-1}(n-1)2^{n-2}$$

- 2. Let f(x, y) be a continuous, real-valued function on \mathbb{R}^2 . Suppose that, for every rectangular region R of area 1, the double integral of f(x, y) over R equals 0. Must f be identically 0?
- 3. Is there a strictly increasing function $f : \mathbb{R} \to \mathbb{R}$ such that f'(x) = f(f(x)) for all x?
- 4. Let P(x) be a polynomial of degree *n* such that P(x) = Q(x)P''(x), where Q(x) is a quadratic polynomial and P''(x) is the second derivative of P(x). Show that if P(x) has at least two distinct roots then it must have *n* distinct roots.
- 5. Determine all positive integers N for which the sphere

$$x^2 + y^2 + z^2 = N$$

has an inscribed regular tetrahedron whose vertices have integer coordinates.

- 6. A class with 2N students took a quiz, on which the possible scores were $0, 1, \ldots, 10$. Each of these scores occurred at least once, and the average score was exactly 7.4. Show that the class can be divided into two groups of N students in such a way that the average score for each group was exactly 7.4.
- 7. Four points are chosen uniformly and independently at random in the interior of a given circle. Find the probability that they are the vertices of a convex quadrilateral.