

Linear Algebra

Book 2

$$f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$f(x, y) = (3x+2y, x-5y)$ can be represented as a matrix transformation

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{bmatrix} 3 & 2 \\ 1 & -5 \end{bmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 3x+2y \\ x-5y \end{pmatrix}$$

Every linear operator can be expressed as matrix multiplication

eg. consider solutions of $y''+y=0$ i.e. $f(x) = \underbrace{a \sin x + b \cos x}_{\begin{pmatrix} a \\ b \end{pmatrix}}$

$$Df(x) = \underbrace{a \cos x - b \sin x}_{\begin{pmatrix} -b \\ a \end{pmatrix}}$$

$$D(rf+sg) = rDf + sDg \quad \begin{pmatrix} -b \\ a \end{pmatrix}$$

$$(rf+sg)' = rf' + sg'$$

$$\underbrace{\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}}_M \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} -b \\ a \end{pmatrix}$$

$$M = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$


$$M^2 = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$

$$M^3 = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

$$M^4 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Every 2×2 real matrix A represents a linear transformation $T_A: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ which is the matrix transformation $T_A \begin{bmatrix} x \\ y \end{bmatrix} = A \begin{bmatrix} x \\ y \end{bmatrix}$.

eg. $\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -y \\ x \end{bmatrix}$ T_A is a counter-clockwise 90° rotation about the origin in \mathbb{R}^2 :




$$T_A \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$T_A \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

$$T_A^{-1} = I \quad I \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ y \end{bmatrix}$$


A counterclockwise rotation by angle θ about the origin in \mathbb{R}^2 represented by the matrix $R_\theta = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \ \theta \end{bmatrix}$ $R_\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}$ $R_\theta \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$




$$\begin{aligned} \cos(\alpha + \beta) &= \cos \alpha \cos \beta - \sin \alpha \sin \beta \\ \sin(\alpha + \beta) &= \sin \alpha \cos \beta + \cos \alpha \sin \beta \end{aligned}$$

$$R_\beta R_\alpha = R_{\alpha + \beta} \quad \begin{bmatrix} \cos \beta & -\sin \beta \\ \sin \beta & \cos \beta \end{bmatrix} \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix} = \begin{bmatrix} \cos(\alpha + \beta) & -\sin(\alpha + \beta) \\ \sin(\alpha + \beta) & \cos(\alpha + \beta) \end{bmatrix}$$


Eg. $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} y \\ x \end{pmatrix}$ is a reflection about the line $y=x$



$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ represents a reflection in the x-axis



$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ represents a shear




Every matrix transformation is a linear transformation: it takes \mathbb{D} to \mathbb{D} and it takes lines to lines. It may distort distances and angles or points.

The null space of a linear transformation $\text{Nul } T = \{ \underline{v} : T\underline{v} = \underline{0} \}$. (the set of null vectors of T)

Recall: $T\underline{0} = \underline{0}$

$$\text{Nul} \begin{bmatrix} 2 & 1 \\ 6 & 3 \end{bmatrix} = \text{Nul } T_A = \left\{ \begin{bmatrix} x \\ -2x \end{bmatrix} : x \in \mathbb{R} \right\}$$

$$A \begin{bmatrix} x \\ -2x \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$



T is one-to-one iff $\text{Nul } T = \{ \underline{0} \}$ (the only null vector is $\underline{0}$).

This statement should be clear:

On the one hand, suppose T is one-to-one.

If $\underline{v} \in \text{Nul } T$ then $T\underline{v} = \underline{0} = T\underline{0}$ then $\underline{v} = \underline{0}$.

This says: if T is one-to-one then $\text{Nul } T = \{ \underline{0} \}$

Conversely, suppose $\text{Nul } T = \{ \underline{0} \}$.

If $T\underline{v} = T\underline{w}$ then $T(\underline{v} - \underline{w}) = T\underline{v} - T\underline{w} = \underline{0}$

so $\underline{v} - \underline{w} \in \text{Nul } T$ i.e. $\underline{v} - \underline{w} = \underline{0}$ i.e. $\underline{v} = \underline{w}$.


"Span" can be used as a noun or as a verb.

The span of a list of vectors v_1, \dots, v_k is the set of all linear combinations of v_1, \dots, v_k .

eg. the span of the vectors $v_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, v_2 = \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix}$ in \mathbb{R}^3 is

the plane $x + y + z = 0$
in \mathbb{R}^3


i.e. the plane $z = -x - y$.



We say that the span of v_1 and v_2 is the plane

or: v_1 and v_2 span the plane $x + y + z = 0$.

eg. the plane $5x + 3y + 7z = 0$ is spanned by $\begin{bmatrix} -3 \\ 5 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 7 \\ 0 \\ -5 \end{bmatrix}$



Given any set of vectors $S \subset \mathbb{R}^3$, the span of S (denoted $\text{span } S = \{\text{linear combinations of vectors in } S\}$) is either $\{0\}$, or a line through 0 , or a plane through 0 , or \mathbb{R}^3 .

Friday: Quiz 5 on Span.

The image of T is $\{T_A \underline{v} : \underline{v} \in \text{domain of } T_A\}$ is the span of the columns of A .

Eg. $A = \begin{bmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{bmatrix}$ defines a linear transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$


(here \mathbb{R}^3 consists of 3×1 column vectors)

$$T_A(v) = A \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} y-z \\ -x+z \\ x-y \end{bmatrix}$$

$$v = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

$$\text{The image of } T_A \text{ is } \left\{ T_A v : v \in \mathbb{R}^3 \right\} = \left\{ \begin{bmatrix} y-z \\ -x+z \\ x-y \end{bmatrix} : x, y, z \in \mathbb{R} \right\}$$

The image of T_A is the span of the columns of A .



$$x \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} + y \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} + z \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$$


(a linear combination of the columns of A)

T_A is not onto \mathbb{R}^3 . This happens because the columns of A fail to span \mathbb{R}^3 .

Any 3 linearly independent vectors in \mathbb{R}^3 will span all of \mathbb{R}^3 (their span is \mathbb{R}^3).

Another example: $B = \begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix}$ defines a linear transformation $T_B: \mathbb{R}^3 \rightarrow \mathbb{R}^3$.


Once again T_B is not onto \mathbb{R}^3 ; its image is the span of the columns of B i.e. the plane $x+y+z=0$ through the origin in \mathbb{R}^3 .



$C = \begin{bmatrix} 3 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix}$ has three linearly independent columns spanning \mathbb{R}^3 i.e. the image of T_C is \mathbb{R}^3 i.e. T_C is onto \mathbb{R}^3 .

Check: If $a \begin{bmatrix} 3 \\ -1 \\ -1 \end{bmatrix} + b \begin{bmatrix} -1 \\ 2 \\ -1 \end{bmatrix} + c \begin{bmatrix} -1 \\ -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 3a-b-c \\ -a+2b-c \\ -a-b+2c \end{bmatrix}$

$A = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$ has $\left\{ \begin{bmatrix} x \\ y \\ 0 \end{bmatrix} : x, y \in \mathbb{R} \right\}$ as the span of its columns.
 T_A is not onto.



The span of the rows of A is $\{ [a, 2a, b] : a, b \in \mathbb{R} \}$