

Linear Algebra

Book 3

Eg. $A = \begin{bmatrix} 1 & 3 & 0 & 4 \\ 2 & 4 & 11 & 7 \\ 0 & 3 & 0 & 4 \\ 1 & 6 & 3 & 5 \end{bmatrix}$

Expanding along the third row, $\det A = 0 \begin{vmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{vmatrix} - 3 \begin{vmatrix} 1 & 0 & 4 \\ 2 & 11 & 7 \\ 1 & 3 & 5 \end{vmatrix} + 0 \begin{vmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{vmatrix} - 4 \begin{vmatrix} 1 & 3 & 0 \\ 2 & 4 & 11 \\ 1 & 6 & 3 \end{vmatrix}$

$$= -3 \left(\begin{vmatrix} 11 & 7 \\ 3 & 5 \end{vmatrix} + 4 \begin{vmatrix} 2 & 11 \\ 1 & 3 \end{vmatrix} \right) - 4 \left(\begin{vmatrix} 1 & 11 \\ 6 & 3 \end{vmatrix} - 3 \begin{vmatrix} 2 & 11 \\ 1 & 3 \end{vmatrix} \right)$$

$$= -3(55 - 21 + 4(6 - 11)) - 4(12 - 66 - 3(6 - 11))$$

$$= 669.$$

(I checked this by computer.)

Wed. Nov. 8 Test. Come a few minutes early if you can.

No Quiz Fri. Nov. 10, 17.

I am away Fri. Nov. 17, Mon. Nov. 20. Lectures for those two days will be prerecorded - check the websites.

Recall: if $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ then $\det A = ad - bc$. A is invertible iff $\det A \neq 0$, in which case $A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$.

This formula has a generalization for $n \times n$ matrices (Cramer's Rule). This is useful although not the most computationally efficient way to compute A^{-1} if n is large.

On HW 2 you had to find A^{-1} where A is 4×4 . The entries of A^{-1} have a common denominator $\det A$.

Eg. $A = \begin{bmatrix} 1 & 2 & 5 \\ 2 & 1 & 3 \\ 7 & 6 & 4 \end{bmatrix}$, $\det A = \begin{vmatrix} 1 & 2 & 5 \\ 2 & 1 & 3 \\ 7 & 6 & 4 \end{vmatrix} = \begin{vmatrix} 1 & 2 & 5 \\ 0 & -3 & -7 \\ 7 & 6 & 4 \end{vmatrix} = - \begin{vmatrix} 1 & 2 & 5 \\ 0 & 3 & 7 \\ 7 & 6 & 4 \end{vmatrix} = - \begin{vmatrix} 1 & 2 & 5 \\ 0 & 3 & 7 \\ 0 & -8 & -31 \end{vmatrix} = \begin{vmatrix} 1 & 2 & 5 \\ 0 & 3 & 7 \\ 0 & -1 & 10 \end{vmatrix}$

$$= |1| \begin{vmatrix} 3 & 7 \\ -1 & 10 \end{vmatrix} = 1 \cdot 37.$$

A^{-1} has fractional entries with common denominator 37.

Matrix of minors: $M = \begin{bmatrix} \begin{vmatrix} 1 & 3 \\ 2 & 4 \end{vmatrix} & \begin{vmatrix} 2 & 5 \\ 7 & 6 \end{vmatrix} & \begin{vmatrix} 2 & 1 \\ 7 & 6 \end{vmatrix} \\ \begin{vmatrix} 2 & 5 \\ 6 & 4 \end{vmatrix} & \begin{vmatrix} 1 & 5 \\ 7 & 4 \end{vmatrix} & \begin{vmatrix} 1 & 2 \\ 7 & 6 \end{vmatrix} \\ \begin{vmatrix} 2 & 5 \\ 1 & 3 \end{vmatrix} & \begin{vmatrix} 1 & 5 \\ 2 & 3 \end{vmatrix} & \begin{vmatrix} 1 & 2 \\ 1 & 1 \end{vmatrix} \end{bmatrix} = \begin{bmatrix} -14 & -13 & 5 \\ -22 & -31 & -8 \\ 1 & -7 & -3 \end{bmatrix}$

$$A^{-1} = \frac{1}{37} \begin{bmatrix} -14 & 22 & 1 \\ 13 & -31 & 7 \\ 5 & 8 & -3 \end{bmatrix} = \begin{bmatrix} -\frac{14}{37} & \frac{22}{37} & \frac{1}{37} \\ \frac{13}{37} & -\frac{31}{37} & \frac{7}{37} \\ \frac{5}{37} & \frac{8}{37} & -\frac{3}{37} \end{bmatrix}$$

← transpose;
apply checkerboard;
divide by det A

Check: $A A^{-1} = \frac{1}{37} \begin{bmatrix} 1 & 2 & 5 \\ 2 & 1 & 3 \\ 7 & 6 & 4 \end{bmatrix} \begin{bmatrix} -14 & 22 & 1 \\ 13 & -31 & 7 \\ 5 & 8 & -3 \end{bmatrix} = \frac{1}{37} \begin{bmatrix} 37 & 0 & 0 \\ 0 & 37 & 0 \\ 0 & 0 & 37 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \checkmark$

If A is a square matrix with integer entries and $\det A = \pm 1$, then A^{-1} also has integer entries.

Find a constant c such that the following matrix has determinant zero:

$$A = \begin{bmatrix} 5 & 3 & 6 \\ 1 & 2 & 4 \\ 7 & 7 & c \end{bmatrix} \begin{array}{l} \leftarrow u = (5 \ 3 \ 6) \\ \leftarrow v = (1 \ 2 \ 4) \\ \leftarrow w \quad u + 2v = (7 \ 7 \ 14) \end{array}$$


If $c=14$ then A has linearly dependent rows so $\det A = 0$ in this case (A is not invertible).

If $c \neq 14$ then A has linearly independent rows then $w \neq (7 \ 7 \ 14)$ and $(0 \ 0 \ 1)$ is a linear combination of u, v, w i.e. Row A contains $u, v, (0 \ 0 \ 1)$.

$$\det \begin{bmatrix} 5 & 3 & 6 \\ 1 & 2 & 4 \\ 0 & 0 & 1 \end{bmatrix} = 7 \times 1 = 7 \neq 0$$


If $A = \begin{bmatrix} -25 & 36 \\ -18 & 26 \end{bmatrix}$, then $A^{10} =$

If $D = \begin{bmatrix} -1 & 0 \\ 0 & 2 \end{bmatrix}$, then $D^{10} =$



$$\det A = -2$$

$$\begin{vmatrix} -25 & 36 \\ -18 & 26 \end{vmatrix} = -25 \times 26 + 36 \times 18 = -2$$



Basis $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ standard basis
 $\begin{bmatrix} x \\ y \end{bmatrix} = xe_1 + ye_2$

There is a basis $\{u, v\}$ for \mathbb{R}^2 such that $Au = -u$, $Av = 2v$

$$A^{10}u = AAA \dots Av$$

$$A^2u = AAu = A(-u) = -Au = u$$

$$A^3u = AAAu = -u$$

$$\vdots$$

$$A^{10}u = u$$

$$A^2v = AA v = A(2v) = 2Av = 4v$$

$$A^3v = 8v$$

$$A^{10}v = \frac{1024}{2}v$$

u, v are eigen vectors of A with corresponding eigenvalues $-1, 2$.

Definition If A is an $n \times n$ matrix, and $v \in \mathbb{R}^n$, then v is an eigenvector for A with eigenvalue λ if

$$Av = \lambda v.$$

How do we find eigenvalues and eigenvectors?

If $Av = \lambda v$ then $Av - \lambda v = 0$ i.e. $Av - \lambda Iv = 0$ i.e. $(A - \lambda I)v = 0$.

We should assume $v \neq 0$ is a nonzero null vector for $A - \lambda I$. This can only happen if $\det(A - \lambda I) = 0$.

This condition allows us to solve for the corresponding eigenvalue λ . Solve for λ ; and for each value λ (each eigenvalue), solve $(A - \lambda I)v = 0$ for the corresponding eigenvector(s) v .

$$\text{For } A = \begin{bmatrix} 25 & 36 \\ -18 & 26 \end{bmatrix}, \quad A - \lambda I = \begin{bmatrix} 25 & 36 \\ -18 & 26 \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} = \begin{bmatrix} 25-\lambda & 36 \\ -18 & 26-\lambda \end{bmatrix}.$$

$$\begin{vmatrix} 25-\lambda & 36 \\ -18 & 26-\lambda \end{vmatrix} = (25-\lambda)(26-\lambda) + 36 \cdot 18 = \lambda^2 - \lambda - 2 = (\lambda+1)(\lambda-2)$$