


# Linear Algebra

Book 1

Example: Find all  $(x, y)$  such that  $5x + 3y = 25$  and  $2x - 7y = -31$ .



We are asking for the simultaneous solution of a system of two equations in two unknowns  $x$  and  $y$ .

$$\begin{cases} 5x + 3y = 25 & (1) \\ 2x - 7y = -31 & (2) \end{cases}$$

$$\begin{aligned} 11y &= 205 \\ y &= 5 \end{aligned}$$

$$\begin{aligned} 5x + 15 &= 25 \\ 5x &= 10 \\ x &= 2 \end{aligned}$$


$$\begin{aligned} 2 \times (1) - 5 \times (2) &= (3) \\ (1) &= (3) \div 4 \end{aligned}$$

$$2 \times 3 - 5(-7) = 6 + 35 = 41$$

$$2 \times 25 - 5 \times (-31) = 50 + 155 = 205$$

Solution:  $(x, y) = (2, 5)$  is the unique solution.

Example: Find all  $(x, y)$  such that  $5x + 3y = 25$  and  $10x + 6y = 17$ .




This system is inconsistent: it has no solution.

$$\begin{cases} 5x + 3y = 25 & (1) \\ 10x + 6y = 17 & (2) \end{cases}$$

$$0 = 33 \quad 2 \times (1) - (2)$$

This is inconsistent.

Example: Find all  $(x, y)$  such that  $5x + 3y = 25$  and  $15x + 9y = 75$ .



This system is consistent but the solution is not unique: there are infinitely many solutions.

$$\begin{array}{rcl} 5x + 3y = 25 & (1) \\ 15x + 9y = 75 & (2) \\ \hline 0 = 0 & (3) = 3 \times (1) - (2) \end{array}$$

$$5x + 3y = 25$$
$$15x + 9y = 75$$

A system of  $m$  linear equations in  $n$  unknowns has the form

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2 \\ \dots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m \end{cases}$$

$(a_{ij}, b_i$  constants for  $i \in \{1, \dots, m\}$ ,  $j \in \{1, 2, \dots, n\}$ ;  $x_1, \dots, x_n$  variables representing unknowns).

Typically, when  $m = n$  we can expect a unique solution;  
 $m > n$  : no solution (inconsistent system);  
 $m < n$  : more than one solution.

Example with  $m=n=3$ : a system of 3 linear equations in 3 unknowns.  
Kim buys a bag of 26 items weighing 226 oz. costing \$34. The items included

cans of tuna (\$1 each, 5oz each)

apples (\$1 each, 8oz each)

loaves of bread (\$3 each, 20oz each)

How many of each item did Kim buy? (say  $x$  cans of tuna,  $y$  apples,  $z$  loaves of bread)

$$x + y + z = 26 \quad (1)$$

$$5x + 8y + 20z = 226 \quad (2)$$

$$x + y + 3z = 34 \quad (3)$$

$$2z = 8 \quad (3) - (1) = (4)$$

$$z = 4 \quad (5)$$

$$x + y = 22 \quad (6) = (8) - (5)$$

$$5x + 8y = 146 \quad (7)$$

$$3y = 36 \quad (7) - 5 \times (6) = (8)$$

$$y = 12 \quad (9) = (8) \div 3$$

$$x = 10 \quad (10) = (6) - (9)$$

$$146 - 5 \times 22 = 146 - 110 = 36$$

The unique solution of this system is  $(x, y, z) = (10, 12, 4)$ .

(Kim bought 10 cans of tuna, 12 apples, and 4 loaves of bread.)

Check! that all three equations are satisfied.

# Matrix formulation of linear systems

$$\begin{aligned} x + y + z &= 26 \\ 5x + 8y + 20z &= 226 \\ x + y + 3z &= 34 \end{aligned} \quad \rightarrow \quad \begin{array}{ccc|c} x & y & z & \text{total} \\ \hline 1 & 1 & 1 & 26 \\ 5 & 8 & 20 & 226 \\ 1 & 1 & 3 & 34 \end{array}$$

$$\begin{aligned} \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 26 \\ 5 & 8 & 20 & 226 \\ 1 & 1 & 3 & 34 \end{array} \right] &\sim \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 26 \\ 5 & 8 & 20 & 226 \\ 0 & 0 & 2 & 8 \end{array} \right] &\sim \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 26 \\ 5 & 8 & 20 & 226 \\ 0 & 0 & 1 & 4 \end{array} \right] &\sim \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 26 \\ 0 & 3 & 15 & 16 \\ 0 & 0 & 1 & 4 \end{array} \right] &\sim \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 26 \\ 0 & 1 & 5 & 32 \\ 0 & 0 & 1 & 4 \end{array} \right] \end{aligned}$$

subtract row 1 from row 3
divide row 3 by 2
subtract 5 times row 1 from row 2
divide row 2 by 3

$$226 - 5 \times 26 = 226 - 130 = 96$$

$$\sim \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 26 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 4 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 0 & 1 & 14 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 4 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 10 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 4 \end{array} \right] \quad \text{i.e.} \quad \begin{aligned} x &= 10 \\ y &= 12 \\ z &= 4 \end{aligned}$$

subtract 5 times row 3 from row 2
subtract row 2 from row 1
subtract row 3 from row 1

Example: Find all  $(x, y)$  such that  $5x + 3y = 25$  and  $2x - 7y = -31$ .

$$\begin{aligned} \left[ \begin{array}{cc|c} 5 & 3 & 25 \\ 2 & -7 & -31 \end{array} \right] &\sim \left[ \begin{array}{cc|c} 1 & \frac{3}{5} & 5 \\ 2 & -7 & -31 \end{array} \right] &\sim \left[ \begin{array}{cc|c} 1 & \frac{3}{5} & 5 \\ 0 & -\frac{41}{5} & -41 \end{array} \right] &\sim \left[ \begin{array}{cc|c} 1 & \frac{3}{5} & 5 \\ 0 & 1 & 5 \end{array} \right] &\sim \left[ \begin{array}{cc|c} 1 & 0 & 2 \\ 0 & 1 & 5 \end{array} \right] \end{aligned}$$

divide row 1 by 5
subtract 2 times row 1 from row 2
multiply row 2 by  $-\frac{5}{41}$ 
subtract  $\frac{3}{5}$  times row 2 from row 1

$$-7 - \frac{6}{5} = -\frac{35}{5} - \frac{6}{5} = -\frac{41}{5}$$

$$-31 - 10 = -41$$

Solution:  $(x, y) = (2, 5)$ .

Alternatively:

$$\left[ \begin{array}{cc|c} 5 & 3 & 25 \\ 2 & -7 & -31 \end{array} \right] \sim \left[ \begin{array}{cc|c} 2 & -7 & -31 \\ 5 & 3 & 25 \end{array} \right] \sim \left[ \begin{array}{cc|c} 1 & -\frac{7}{2} & -\frac{31}{2} \\ 5 & 3 & 25 \end{array} \right] \sim \dots \sim \left[ \begin{array}{cc|c} 1 & 0 & 2 \\ 0 & 1 & 5 \end{array} \right]$$

interchange rows 1 and 2

Even better:  $\begin{bmatrix} 5 & 3 & 25 \\ 2 & -7 & -31 \end{bmatrix} \sim \begin{bmatrix} 1 & 17 & 87 \\ 2 & -7 & -31 \end{bmatrix} \sim \begin{bmatrix} 1 & 17 & 87 \\ 0 & -41 & -205 \end{bmatrix} \sim \begin{bmatrix} 1 & 17 & 87 \\ 0 & 1 & 5 \end{bmatrix}$

subtract 2 times row 2 from row 1

subtract 2 times row 1 from row 2

divide row 2 by -41

$$\begin{aligned} & -31 - 2 \times 87 \\ & = -31 - 174 \\ & = -205 \end{aligned}$$

$$\sim \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & 5 \end{bmatrix}$$

Solution:  $(x, y) = (2, 5)$ .

subtract 17 times row 2 from row 1

check!

$$\begin{aligned} 5 \times 2 + 3 \times 5 &= 25 \\ 2 \times 2 - 7 \times 5 &= -31 \end{aligned}$$

Elementary row operations:

- (i) add a multiple of one row to another
- (ii) multiply a row by a nonzero constant
- (iii) interchange two rows

$A \sim B$  means that  $A, B$  are linear systems having the same solutions.

We use Gaussian elimination to reduce  $A_1 \sim A_2 \sim \dots \sim A_n$  where  $A_1$  represents the linear system and  $A_n$  represents an equivalent linear system (i.e. having the same solutions) but  $A_n$  is simpler than  $A_1$ . Each step  $A_i \sim A_{i+1}$  is obtained by one elementary row operation.

Why just one operation at a time?

$$\begin{cases} 5x + 3y = 25 \\ 2x - 7y = -31 \end{cases}$$

$$\begin{bmatrix} 5 & 3 & 25 \\ 2 & -7 & -31 \end{bmatrix} \sim \begin{bmatrix} 1 & \frac{3}{5} & 5 \\ 2 & -7 & -31 \end{bmatrix}$$

divide row 1 by 5

divide row 2 by 2

$$\begin{bmatrix} 1 & \frac{3}{5} & 5 \\ 0 & -\frac{21}{5} & -\frac{41}{5} \end{bmatrix} \sim \begin{bmatrix} 1 & \frac{3}{5} & 5 \\ 0 & 1 & 5 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 5 \\ 0 & 1 & 5 \end{bmatrix}$$

subtract row 2 from row 1

subtract row 1 from row 2

i.e.  $y = 5$

$0 = 0$



unique solution  $(2, 5)$

infinitely many solutions

Gauss



Gaussian distribution

$\begin{bmatrix} 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$ ,  $\begin{bmatrix} 1 & 0 & 7 \\ 0 & 0 & 9 \end{bmatrix}$ ,  $\begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & 5 \end{bmatrix}$  are examples of matrices in reduced row echelon form:  
they cannot be simplified any further by elementary row operations.

$\begin{bmatrix} 1 & 17 & 87 \\ 0 & 1 & 5 \end{bmatrix}$  is almost reduced; it is in row echelon form.

For a linear system whose matrix is in row echelon form, we can solve for the unknowns  $x_1, x_2, \dots, x_n$ , we solve for  $x_n$ , then  $x_{n-1}$ , then  $x_{n-2}, \dots, x_1$  by back-substitution.

eg.  $\begin{bmatrix} 5 & 3 & 7 & 3 \\ 0 & 2 & 11 & 4 \\ 0 & 0 & 6 & 8 \end{bmatrix}$  is in row echelon form.

Every linear system has a unique reduced row echelon form.

In any  $m \times n$  matrix, a pivot is the first nonzero entry in its row.  
(Pivots are highlighted above.)

In order for a matrix to be in row echelon form, we must have

- pivots in any row must occur to the right of pivots in any previous rows;
- any zero rows occur at the bottom.

Assuming a matrix is already in row echelon form, then to be in reduced row echelon form, we must have

- every pivot entry must be a 1
- every column having a pivot has only one nonzero entry.

Example: Solve the following linear system of 3 equations in 5 unknowns:

$$\begin{cases} x_1 + 4x_2 - x_3 + 2x_4 + 3x_5 = 6 \\ 2x_1 + 8x_2 - x_3 + 7x_4 + 4x_5 = 19 \\ -x_1 - 4x_4 + 4x_3 + 8x_4 - 4x_5 = 26 \end{cases}$$

$$\left[ \begin{array}{ccccc|c} 1 & 4 & -1 & 2 & 3 & 6 \\ 2 & 8 & -1 & 7 & 4 & 19 \\ -1 & -4 & 4 & 8 & -4 & 26 \end{array} \right] \sim \left[ \begin{array}{ccccc|c} 1 & 4 & -1 & 2 & 3 & 6 \\ 0 & 0 & 1 & 3 & -2 & 7 \\ -1 & -4 & 4 & 8 & -4 & 26 \end{array} \right] \sim \left[ \begin{array}{ccccc|c} 1 & 4 & -1 & 2 & 3 & 6 \\ 0 & 0 & 1 & 3 & -2 & 7 \\ 0 & 0 & 3 & 10 & -1 & 32 \end{array} \right]$$

$$\sim \left[ \begin{array}{ccccc|c} 1 & 4 & -1 & 2 & 3 & 6 \\ 0 & 0 & 1 & 3 & -2 & 7 \\ 0 & 0 & 0 & 5 & 5 & 11 \end{array} \right]$$

This matrix is in row echelon form. This can be used to solve the linear system by back-substitution.

$$x_4 + 5x_5 = 11 \quad x_5 = t \text{ is a free parameter.}$$

$$x_4 = 11 - 5t$$

$$x_3 + 3x_4 - 2x_5 = 7$$

$$x_3 = 7 - 3x_4 + 2x_5 = 7 - 3(11 - 5t) + 2t = -26 + 17t$$

$$x_1 + 4x_2 - x_3 + 2x_4 + 3x_5 = 6$$


$$x_2 = s \text{ is another free parameter}$$

$$x_1 = 6 - 4x_2 + x_3 - 2x_4 - 3x_5 = 6 - 4s + (-26 + 17t) - 2(11 - 5t) - 3t = -42 - 4s + 24t$$

Solutions:  $(x_1, x_2, x_3, x_4, x_5) = (-42 - 4s + 24t, s, -26 + 17t, 11 - 5t, t)$  where  $s, t$  are arbitrary.

Geometrically, the set of solutions forms a plane (2-dimensional surface) in  $\mathbb{R}^5$ .

two parameters  $s, t$  are coordinates for the plane



Solution set inside  $\mathbb{R}^5$ .

The point corresponding to  $(s, t) = (3, 1)$  is  $(30, 3, -9, 6, 1)$  is another solution.

Our system is consistent but the solution is not unique.



$$\left[ \begin{array}{cccc|c} 1 & 4 & -1 & 2 & 3 & 6 \\ 0 & 0 & 1 & 3 & -2 & 7 \\ 0 & 0 & 0 & 1 & 5 & 11 \end{array} \right] \sim \left[ \begin{array}{cccc|c} 1 & 4 & 0 & 5 & 1 & 13 \\ 0 & 0 & 1 & 3 & -2 & 7 \\ 0 & 0 & 0 & 1 & 5 & 11 \end{array} \right] \sim \left[ \begin{array}{cccc|c} 1 & 4 & 0 & 0 & -24 & -42 \\ 0 & 0 & 1 & 3 & -2 & 7 \\ 0 & 0 & 0 & 1 & 5 & 11 \end{array} \right] \sim \left[ \begin{array}{cccc|c} 1 & 4 & 0 & 0 & -24 & -42 \\ 0 & 0 & 1 & 0 & -17 & -26 \\ 0 & 0 & 0 & 1 & 5 & 11 \end{array} \right]$$

(row echelon form) (reduced row echelon form)

To solve a linear system in reduced row echelon form, introduce parameters for the free variables (the variables whose columns do not contain a pivot).

In the example above,  $x_2$  and  $x_5$  are the free variables. Introduce  $s, t$ .  $x_2 = s, x_5 = t$  can be chosen freely. Solve for the variables  $x_1, x_3, x_4$  using the equations appearing in the reduced row echelon form:

$$\left. \begin{array}{l} x_1 + 4s - 2t = -42 \\ x_3 - 17t = -26 \\ x_4 + 5t = 11 \end{array} \right\} \Rightarrow (x_1, x_2, x_3, x_4, x_5) = (-42 - 4s + 2t, s, -26 + 17t, 11 - 5t, t)$$

where  $s, t$  are arbitrary parameters. (This is the parametric solution in terms of the parameters  $s, t$ . The system is consistent, having infinitely many solutions.)

As long as the rightmost column has no pivot, the system is consistent.

The general solution can be written as

$$\begin{aligned} (x_1, x_2, x_3, x_4, x_5) &= (-42 - 4s + 2t, s, -26 + 17t, 11 - 5t, t) \\ &= (-42, 0, -26, 11, 0) + s(-4, 1, 0, 0, 0) + t(2, 0, 17, -5, 1) \end{aligned}$$

$$(a_1, a_2, \dots, a_n) + (b_1, b_2, \dots, b_n) = (a_1 + b_1, a_2 + b_2, \dots, a_n + b_n) \quad (\text{vector addition})$$

$$c(a_1, a_2, \dots, a_n) = (ca_1, ca_2, \dots, ca_n) \quad (\text{scalar multiplication})$$

↑ scalar      ↑ vector

# Algebraic operations for matrices

If  $A = \begin{bmatrix} 2 & 3 & 5 \\ 1 & -7 & 11 \end{bmatrix}$  and  $B = \begin{bmatrix} 6 & 1 \\ -2 & 3 \end{bmatrix}$

then  $BA = \begin{bmatrix} 6 & 1 \\ -2 & 3 \end{bmatrix} \begin{bmatrix} 2 & 3 & 5 \\ 1 & -7 & 11 \end{bmatrix} = \begin{bmatrix} 13 & 11 & 41 \\ -1 & -27 & 23 \end{bmatrix}$

$2 \times 2$        $2 \times 3$        $2 \times 3$

Here  $AB$  is undefined.

An  $m \times n$  matrix has the form

$$A = \begin{bmatrix} a_{1,1} & a_{1,2} & \dots & a_{1,n} \\ a_{2,1} & a_{2,2} & \dots & a_{2,n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m,1} & a_{m,2} & \dots & a_{m,n} \end{bmatrix}$$

$a_{i,j}$  is the  $(i,j)$ -entry of the matrix  $A$ .

$$i \in \{1, 2, \dots, m\}$$

$$j \in \{1, 2, \dots, n\}$$

Often  $a_{i,j}$  is written  $a_{ij}$ .  
(unless this results in confusion)

If  $A$  is  $m \times n$  and  $B$  is  $n \times r$  then  $AB$  is  $m \times r$ .

We can't multiply two matrices unless the number of columns in the first matrix equals the number of rows in the second matrix.

eg.  $\begin{bmatrix} 2 & 3 & 5 \\ 1 & -7 & 11 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} 12 & 8 \\ 23 & 4 \end{bmatrix}$  whereas  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 2 & 3 & 5 \\ 1 & -7 & 11 \end{bmatrix} = \begin{bmatrix} 2 & 3 & 5 \\ 1 & -7 & 11 \\ 5 & -1 & 21 \end{bmatrix}$

$2 \times 3$        $3 \times 2$        $3 \times 2$        $2 \times 3$