

# Linear Algebra

Book 3

Eg.  $A = \begin{bmatrix} 1 & 3 & 0 & 4 \\ 2 & 4 & 11 & 7 \\ 0 & 3 & 0 & 4 \\ 1 & 6 & 3 & 5 \end{bmatrix}$

Expanding along the third row,  $\det A = 0 \begin{vmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{vmatrix} - 3 \begin{vmatrix} 1 & 0 & 4 \\ 2 & 11 & 7 \\ 1 & 3 & 5 \end{vmatrix} + 0 \begin{vmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{vmatrix} - 4 \begin{vmatrix} 1 & 3 & 0 \\ 2 & 4 & 11 \\ 1 & 6 & 3 \end{vmatrix}$

$$= -3 \left( \begin{vmatrix} 11 & 7 \\ 3 & 5 \end{vmatrix} + 4 \begin{vmatrix} 2 & 11 \\ 1 & 3 \end{vmatrix} \right) - 4 \left( \begin{vmatrix} 1 & 11 \\ 6 & 3 \end{vmatrix} - 3 \begin{vmatrix} 2 & 11 \\ 1 & 3 \end{vmatrix} \right)$$

$$= -3(55 - 21 + 4(6 - 11)) - 4(12 - 66 - 3(6 - 11))$$

$$= 669.$$

(I checked this by computer.)

Wed. Nov. 8 Test. Come a few minutes early if you can.

No Quiz Fri. Nov. 10, 17.

I am away Fri. Nov. 17, Mon. Nov. 20. Lectures for those two days will be prerecorded - check the websites.

Recall: if  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$  then  $\det A = ad - bc$ .  $A$  is invertible iff  $\det A \neq 0$ , in which case  $A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$ .

This formula has a generalization for  $n \times n$  matrices (Cramer's Rule). This is useful although not the most computationally efficient way to compute  $A^{-1}$  if  $n$  is large.

On HW 2 you had to find  $A^{-1}$  where  $A$  is  $4 \times 4$ . The entries of  $A^{-1}$  have a common denominator  $\det A$ .

Eg.  $A = \begin{bmatrix} 1 & 2 & 5 \\ 2 & 1 & 3 \\ 7 & 6 & 4 \end{bmatrix}$ ,  $\det A = \begin{vmatrix} 1 & 2 & 5 \\ 2 & 1 & 3 \\ 7 & 6 & 4 \end{vmatrix} = \begin{vmatrix} 1 & 2 & 5 \\ 0 & -3 & -7 \\ 7 & 6 & 4 \end{vmatrix} = - \begin{vmatrix} 1 & 2 & 5 \\ 0 & 3 & 7 \\ 7 & 6 & 4 \end{vmatrix} = - \begin{vmatrix} 1 & 2 & 5 \\ 0 & 3 & 7 \\ 0 & -8 & -31 \end{vmatrix} = \begin{vmatrix} 1 & 2 & 5 \\ 0 & 3 & 7 \\ 0 & -8 & -31 \end{vmatrix} = \begin{vmatrix} 1 & 2 & 5 \\ 0 & 3 & 7 \\ 0 & -1 & 10 \end{vmatrix}$

$$= |1| \begin{vmatrix} 3 & 7 \\ -1 & 10 \end{vmatrix} = 1 \cdot 37.$$

$A^{-1}$  has fractional entries with common denominator 37.  
 Matrix of minors:  $M = \begin{bmatrix} \begin{vmatrix} 1 & 3 \\ 2 & 4 \end{vmatrix} & \begin{vmatrix} 1 & 5 \\ 2 & 3 \end{vmatrix} & \begin{vmatrix} 2 & 1 \\ 7 & 6 \end{vmatrix} \\ \begin{vmatrix} 2 & 5 \\ 7 & 4 \end{vmatrix} & \begin{vmatrix} 1 & 5 \\ 7 & 4 \end{vmatrix} & \begin{vmatrix} 1 & 2 \\ 7 & 6 \end{vmatrix} \\ \begin{vmatrix} 2 & 5 \\ 1 & 3 \end{vmatrix} & \begin{vmatrix} 1 & 5 \\ 2 & 3 \end{vmatrix} & \begin{vmatrix} 1 & 2 \\ 1 & 1 \end{vmatrix} \end{bmatrix} = \begin{bmatrix} -14 & -13 & 5 \\ -22 & -31 & -8 \\ 1 & -7 & -3 \end{bmatrix}$

$$A^{-1} = \frac{1}{37} \begin{bmatrix} -14 & 22 & 1 \\ 13 & -31 & 7 \\ 5 & 8 & -3 \end{bmatrix} = \begin{bmatrix} -\frac{14}{37} & \frac{22}{37} & \frac{1}{37} \\ \frac{13}{37} & -\frac{31}{37} & \frac{7}{37} \\ \frac{5}{37} & \frac{8}{37} & -\frac{3}{37} \end{bmatrix}$$

← transpose;  
 apply checkerboard;  
 divide by det A

Check:  $A A^{-1} = \frac{1}{37} \begin{bmatrix} 1 & 2 & 5 \\ 2 & 1 & 3 \\ 7 & 6 & 4 \end{bmatrix} \begin{bmatrix} -14 & 22 & 1 \\ 13 & -31 & 7 \\ 5 & 8 & -3 \end{bmatrix} = \frac{1}{37} \begin{bmatrix} 37 & 0 & 0 \\ 0 & 37 & 0 \\ 0 & 0 & 37 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \checkmark$

If  $A$  is a square matrix with integer entries and  $\det A = \pm 1$ , then  $A^{-1}$  also has integer entries.

Find a constant  $c$  such that the following matrix has determinant zero:

$$A = \begin{bmatrix} 5 & 3 & 6 \\ 1 & 2 & 4 \\ 7 & 7 & c \end{bmatrix} \begin{array}{l} \leftarrow u = (5 \ 3 \ 6) \\ \leftarrow v = (1 \ 2 \ 4) \\ \leftarrow w \quad u + 2v = (7 \ 7 \ 14) \end{array}$$


If  $c = 14$  then  $A$  has linearly dependent rows so  $\det A = 0$  in this case ( $A$  is not invertible).

If  $c \neq 14$  then  $A$  has linearly independent rows then  $w \neq (7 \ 7 \ 14)$  and  $(0 \ 0 \ 1)$  is a linear combination of  $u, v, w$  i.e. Row  $A$  contains  $u, v, (0 \ 0 \ 1)$ .

$$\det \begin{bmatrix} 5 & 3 & 6 \\ 1 & 2 & 4 \\ 0 & 0 & 1 \end{bmatrix} = 7 \times 1 = 7 \neq 0.$$


If  $A = \begin{bmatrix} -25 & 36 \\ -18 & 26 \end{bmatrix}$ , then  $A^{10} =$

If  $D = \begin{bmatrix} -1 & 0 \\ 0 & 2 \end{bmatrix}$ , then  $D^{10} = \begin{bmatrix} 1 & 0 \\ 0 & 2^{10} \end{bmatrix}$



$$\det A = -2$$

$$\begin{vmatrix} -25 & 36 \\ -18 & 26 \end{vmatrix} = -25 \times 26 + 36 \times 18 = -2$$



Basis  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$  standard basis  
 $\begin{bmatrix} x \\ y \end{bmatrix} = xe_1 + ye_2$

There is a basis  $\{u, v\}$  for  $\mathbb{R}^2$  such that  $Au = -u$ ,  $Av = 2v$

$$\begin{aligned} A^{10}u &= AAA \dots Av \\ A^2u &= AAu = A(-u) = -Au = u \\ A^3u &= AAAu = -u \\ &\vdots \\ A^{10}u &= u \end{aligned}$$

$$\begin{aligned} A^2v &= AA v = A(2v) = 2Av = 4v \\ A^3v &= 8v \\ A^{10}v &= \frac{1024}{2}v \end{aligned}$$

$u, v$  are eigenvectors of  $A$  with corresponding eigenvalues  $-1, 2$ .

Definition If  $A$  is an  $n \times n$  matrix, and  $v \in \mathbb{R}^n$ , then  $v$  is an eigenvector for  $A$  with eigenvalue  $\lambda$  if

$$Av = \lambda v.$$

How do we find eigenvalues and eigenvectors?

If  $Av = \lambda v$  then  $Av - \lambda v = 0$  i.e.  $(A - \lambda I)v = 0$  i.e.  $(A - \lambda I)v = 0$ .

We should assume  $v \neq 0$  is a nonzero null vector for  $A - \lambda I$ . This can only happen if  $\det(A - \lambda I) = 0$ .

This condition allows us to solve for the corresponding eigenvalue  $\lambda$ . Solve for  $\lambda$ ; and for each value  $\lambda$  (each eigenvalue), solve  $(A - \lambda I)v = 0$  for the corresponding eigenvector(s)  $v$ .

$$\text{For } A = \begin{bmatrix} 25 & 36 \\ -18 & 26 \end{bmatrix}, \quad A - \lambda I = \begin{bmatrix} 25 & 36 \\ -18 & 26 \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} = \begin{bmatrix} 25-\lambda & 36 \\ -18 & 26-\lambda \end{bmatrix}.$$

$$\begin{vmatrix} 25-\lambda & 36 \\ -18 & 26-\lambda \end{vmatrix} = (25-\lambda)(26-\lambda) + 36 \cdot 18 = \lambda^2 - \lambda - 2 = (\lambda+1)(\lambda-2)$$

The characteristic polynomial has two roots  $\lambda_1 = -1$ ,  $\lambda_2 = 2$  (the two eigenvalues).

To find the corresponding eigenvectors  $v_1, v_2$ :

First take  $\lambda_1 = -1$  and solve  $Av_1 = -v_1$  i.e.  $(A+I)v_1 = 0$ .  $A+I = \begin{bmatrix} 24 & 36 \\ -18 & 27 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$  (by inspection)

Or  $\begin{bmatrix} 24 & 36 \\ -18 & 27 \end{bmatrix} \sim \begin{bmatrix} 1 & -\frac{3}{2} \\ 0 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & -\frac{3}{2} \\ 0 & 0 \end{bmatrix}$  has null space  $\text{Span}\left\{\begin{bmatrix} \frac{3}{2} \\ 1 \end{bmatrix}\right\}$  with basis  $\begin{bmatrix} \frac{3}{2} \\ 1 \end{bmatrix}$

$$\begin{bmatrix} 1 & -\frac{3}{2} \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad \therefore \quad x - \frac{3}{2}y = 0 \quad \text{Introduce a parameter } t.$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{3}{2}t \\ t \end{bmatrix} = t \begin{bmatrix} \frac{3}{2} \\ 1 \end{bmatrix}$$

We can take  $v_1$  to be any nonzero scalar multiple of  $\begin{bmatrix} \frac{3}{2} \\ 1 \end{bmatrix}$ . If we take  $v_1 = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$ . So  $Av_1 = \lambda_1 v_1 = -v_1$ .

For  $\lambda_2 = 2$ : Solve  $Av_2 = \lambda_2 v_2 = 2v_2$  i.e.  $(A - 2I)v_2 = 0$  where  $A - 2I = \begin{bmatrix} -25 & 36 \\ -18 & 26 \end{bmatrix} - \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = \begin{bmatrix} -27 & 36 \\ -18 & 24 \end{bmatrix}$

A null vector of  $A - 2I$ :  $v_2 = \begin{bmatrix} 4 \\ 3 \end{bmatrix}$  or  $\begin{bmatrix} 1 \\ 3 \end{bmatrix}$  so  $\begin{bmatrix} -27 & 36 \\ -18 & 24 \end{bmatrix} \begin{bmatrix} 4 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$  i.e.  $Av_2 = \lambda_2 v_2 = 2v_2$ .

$v_1 = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$  and  $v_2 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$  is a basis of  $\mathbb{R}^2$  consisting of eigenvectors of  $A$ .

We started with  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$  as the standard basis.

To find  $A^{10}$ : two approaches.

Let  $B = [v_1 | v_2] = \begin{bmatrix} 3 & 1 \\ 2 & 3 \end{bmatrix}$ . Then  $AB = A \begin{bmatrix} 3 & 1 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} -3 & 8 \\ -2 & 6 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 0 & 2 \end{bmatrix} = BD$ ,  $D = \begin{bmatrix} -1 & 0 \\ 0 & 2 \end{bmatrix}$  (diagonal matrix)

so  $AB B^{-1} = BDB^{-1}$  i.e.  $A = BDB^{-1}$ .

So  $A^{10} = (BDB^{-1})(BDB^{-1}) \dots (BDB^{-1}) = B D^{10} B^{-1} = \begin{bmatrix} 3 & 1 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1024 \end{bmatrix} \begin{bmatrix} 3 & -4 \\ -2 & 3 \end{bmatrix} = \begin{bmatrix} -8183 & 12276 \\ -6138 & 9208 \end{bmatrix}$

To check:  $\det(A^{10}) = (\det A)^{10} = (-2)^{10} = 1024$ .

$\det \begin{bmatrix} \downarrow \\ \end{bmatrix} = 1024$

$\det A = (-25)(26) - (36)(-18) = -2$ .

$\det A = (\det B)(\det D)(\det B^{-1}) = 1 \times (-2) \times 1 = -2$

Second approach:  $A^{10}v_1 = v_1$ ,  $A^{10}v_2 = 1024v_2$

$\left. \begin{aligned} v_1 &= \begin{bmatrix} 3 \\ 2 \end{bmatrix} = 3e_1 + 2e_2 \\ v_2 &= \begin{bmatrix} 1 \\ 3 \end{bmatrix} = 4e_1 + 3e_2 \end{aligned} \right\} \Rightarrow$

$\left. \begin{aligned} e_1 &= 3v_1 - 2v_2 = 3 \begin{bmatrix} 3 \\ 2 \end{bmatrix} - 2 \begin{bmatrix} 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \\ e_2 &= -4v_1 + 3v_2 = -4 \begin{bmatrix} 3 \\ 2 \end{bmatrix} + 3 \begin{bmatrix} 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{aligned} \right\}$

$A^{10}e_1 = A^{10}(3v_1 - 2v_2) = 3 \cdot v_1 - 2 \cdot 1024v_2 = 3 \begin{bmatrix} 3 \\ 2 \end{bmatrix} - 2048 \begin{bmatrix} 1 \\ 3 \end{bmatrix} = \begin{bmatrix} -8183 \\ -6138 \end{bmatrix}$

$A^{10}e_2 = A^{10}(-4v_1 + 3v_2) = -4v_1 + 3 \cdot 1024v_2 = -4 \begin{bmatrix} 3 \\ 2 \end{bmatrix} + 3072 \begin{bmatrix} 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 12276 \\ 9208 \end{bmatrix}$

$A^{10} = \begin{bmatrix} -8183 & 12276 \\ -6138 & 9208 \end{bmatrix}$

$A$  and  $D$  are similar: they represent the same linear transformation with respect to different choices of basis.

Ex. diagonalize the matrix  $A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & 2 \\ 0 & 0 & 3 \end{bmatrix}$ .  $\det A = \begin{vmatrix} 1 & -1 & 1 \\ 2 & 1 & 2 \\ 0 & 0 & 3 \end{vmatrix} = \begin{vmatrix} 1 & -1 \\ 2 & 1 \end{vmatrix} \cdot 3 = 6 \cdot 3 = 18$

First compute the characteristic polynomial  $\det(A - \lambda I) = \begin{vmatrix} 1-\lambda & -1 & 1 \\ 2 & 1-\lambda & 2 \\ 0 & 0 & 3-\lambda \end{vmatrix} = \begin{vmatrix} 1-\lambda & -1 \\ 2 & 1-\lambda \end{vmatrix} (3-\lambda) = [(1-\lambda)(1-\lambda) + 2] (3-\lambda)$   
 $= [\lambda^2 - 5\lambda + 6] (3-\lambda) = (\lambda-2)(\lambda-3)(3-\lambda) = -(\lambda-2)(\lambda-3)^2$  has roots  $2, 3, 3$  (the eigenvalues of  $A$ ).

Find eigenvector  $v_1$  for  $\lambda_1 = 2$ : solve  $(A - \lambda_1 I)v_1 = 0$  i.e.  $\begin{bmatrix} 2 & -1 & 1 \\ 2 & -1 & 2 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ ,  $v_1 = \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} \Rightarrow Av_1 = 2v_1$ .

Find eigenvectors  $v_2, v_3$  for  $\lambda_2 = \lambda_3 = 3$ : solve  $(A - 3I)v = 0$  i.e.  $\begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ . Take  $v_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ ,  $v_3 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$

Note: We want two linearly independent solutions.

Form the matrix  $B = [v_1 | v_2 | v_3] = \begin{bmatrix} 1 & 1 & 0 \\ 2 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$  whose columns are the eigenvectors. ( $v_1, v_2, v_3$  is our basis of eigenvectors)

Then  $AB = BD$  where  $D = \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{bmatrix} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ .


i.e.  $AB B^{-1} = B D B^{-1}$

i.e.  $A = B D B^{-1}$ . We have diagonalized  $A$ .

$AB = A [v_1 | v_2 | v_3] = [Av_1 | Av_2 | Av_3] = [2v_1 | 3v_2 | 3v_3] = \begin{bmatrix} v_1 & v_2 & v_3 \\ \left[ \begin{matrix} 2 & 3 & 3 \end{matrix} \right] \end{bmatrix} = BD$

Check:  $\text{tr} A \stackrel{?}{=} \text{tr} D$ ,  $\det A \stackrel{?}{=} \det D$   
 $8 = 8$ ,  $18 = 18$

$\mathbb{R}^3$  has an eigenvector  $v_1$  with eigenvalue  $\lambda_1 = 2$  and an eigenspace  $\text{Span}\{v_2, v_3\}$  with eigenvalue  $3$ .



The eigenspace for  $\lambda$  is  $\text{Nul}(A - \lambda I) = \{ \text{all eigenvectors having eigenvalue } \lambda \}$   
 $= \{ \text{all } v \text{ satisfying } Av = \lambda v \}$ .

$\begin{bmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{bmatrix}$  has a single eigenspace  $\mathbb{R}^3$  with eigenvalue 5.

Actually, we don't necessarily have a basis of eigenvectors.

Consider  $A = \begin{bmatrix} 7 & 16 \\ -4 & 9 \end{bmatrix}$ .

Find the characteristic polynomial  $\det(A - \lambda I) = \begin{vmatrix} 7-\lambda & 16 \\ -4 & 9-\lambda \end{vmatrix} = (7-\lambda)(9-\lambda) + 64 = \lambda^2 - 2\lambda + 1 = (\lambda-1)^2$   
 which has roots 1, 1. (Only one distinct eigenvalue)  $\text{Tr} A = 2$   $\det A = 1$

Look for eigenvectors:  $(A - I)v = 0$  i.e.  $\begin{bmatrix} 6 & 16 \\ -4 & 8 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$ . Take  $v_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ .

Try to complete this to a basis  $v_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $v_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ;  $B = [v_1 | v_2] = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$

$$AB = A[v_1 | v_2] = [Av_1 | Av_2] = \begin{bmatrix} 2 & 9 \\ 1 & 5 \end{bmatrix} = B \begin{bmatrix} 1 & 4 \\ 0 & 1 \end{bmatrix} = B M$$

$AB = BM \iff A = BMB^{-1}$   
 $A, M$  are similar matrices  
 having the same trace,  
 determinant, characteristic poly.

$$Av_2 = \begin{bmatrix} -7 & 16 \\ -4 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 9 \\ 5 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 4 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 9 \\ 1 & 5 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & ? \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 9 \\ 1 & 5 \end{bmatrix}$$



$$\underbrace{\begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}}_B \underbrace{\begin{bmatrix} 1 & ? \\ 0 & 1 \end{bmatrix}}_M = \underbrace{\begin{bmatrix} 2 & 9 \\ 1 & 5 \end{bmatrix}}_{AB}$$

$$M = B^{-1}AB = \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 2 & 9 \\ 1 & 5 \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ 0 & 1 \end{bmatrix}$$

$$B = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$$

$$B^{-1} = \frac{1}{1} \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix}$$





$A$  is not diagonalizable;  
 $\mathbb{R}^2$  does not have a basis  
 consisting of eigenvectors for  $A$ .  
 (we have one eigenvector only).


An example with no eigenvectors or eigenvalues:

$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$  represents a  $90^\circ$  rotation counterclockwise

Algebraically: compute the characteristic polynomial

$$\det(A - \lambda I) = \det \left( \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} \right) = \det \begin{bmatrix} -\lambda & -1 \\ 1 & -\lambda \end{bmatrix} = \lambda^2 + 1$$

Over  $\mathbb{R}$  there are no roots of  $\lambda^2 + 1$  (you cannot factor this).



Over  $\mathbb{C} = \{a + bi : a, b \in \mathbb{R}\}$ , however, we factor  $\lambda^2 + 1 = (\lambda + i)(\lambda - i)$   
so the roots  $i, -i$  give two eigenvalues in  $\mathbb{C}$ .  $i^2 = -1$

Find eigenvectors for  $A$

$$Av = iv \iff (A - iI)v = 0 \text{ i.e. } \begin{bmatrix} -i & -1 \\ 1 & -i \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}. \text{ Take } v_1 = \begin{bmatrix} i \\ 1 \end{bmatrix} \text{ as an eigenvector.}$$

# Vector Spaces: Chapter 1

Scalars: real numbers / complex numbers / rational numbers / general fields

A field is a set of scalars in which we can add, subtract, multiply and divide.

A vector space is a set  $V$  whose elements are called vectors, including a zero vector  $\underline{0}$ , and operations  $+$ ,  $-$ , scalar multiplication satisfying

1. For  $\underline{u}, \underline{v} \in V$ ,  $\underline{u} + \underline{v} \in V$ . (vector + vector = vector)
2.  $\underline{u} + \underline{v} = \underline{v} + \underline{u}$
3.  $(\underline{u} + \underline{v}) + \underline{w} = \underline{u} + (\underline{v} + \underline{w})$  } for all  $\underline{u}, \underline{v}, \underline{w} \in V$
4.  $\underline{u} + \underline{0} = \underline{u} = \underline{0} + \underline{u}$
5. For each  $\underline{u} \in V$ , there is a vector  $-\underline{u} \in V$  such that  $\underline{u} + (-\underline{u}) = \underline{0}$
6. Scalar multiplication: For every scalar  $c$  and  $\underline{u} \in V$ ,  $c\underline{u} \in V$
7. Distributivity:  $c(\underline{u} + \underline{v}) = c\underline{u} + c\underline{v}$
8.  $\dots$   $(c+d)\underline{u} = c\underline{u} + d\underline{u}$
9. Associativity:  $(cd)\underline{u} = c(d\underline{u})$
10.  $1\underline{u} = \underline{u}$

(scalar + scalar = scalar, ~~scalar + vector~~)

~~vector x vector~~

(scalar x vector = vector)

$$\begin{array}{ccc} & \underline{0} & \underline{u} \\ & \uparrow & \uparrow \\ \text{scalar} & \cdot & \text{vector} \end{array}$$

as follows from the axioms:  $\underline{0}\underline{u} + \underline{0}\underline{u} = (0+0)\underline{u} = \underline{0}\underline{u}$ . Add  $-\underline{0}\underline{u}$  to both sides:

$$(\underline{0}\underline{u} + \underline{0}\underline{u}) + (-\underline{0}\underline{u}) = \underline{0}\underline{u} + (-\underline{0}\underline{u}) = \underline{0}$$

By (3),  $\underline{0}\underline{u} + (\underline{0}\underline{u} + (-\underline{0}\underline{u})) = \underline{0}$

By (5)  $\underline{0}\underline{u} + \underline{0} = \underline{0}$

$$\underline{0}\underline{u} = \underline{0}$$

Examples of vector spaces:

$\mathbb{R}^n$  (actually,  $\mathbb{R}^{n \times 1}$  is column vectors of length  $n$ ;  $\mathbb{R}^{1 \times n}$  is row vectors of length  $n$ ).

Subspaces of  $\mathbb{R}^n$

The set of all polynomials of degree  $< n$  in  $x$  is an  $n$ -dimensional vector space

$$V = \{ a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1} : a_0, a_1, a_2, \dots, a_{n-1} \text{ are scalars} \}$$

$\{ 1, x, x^2, \dots, x^{n-1} \}$  is a basis for  $V$ .  $x$  is an indeterminate (i.e. not a number, just a symbol).

$\{ 1, x, x(x-1), x(x-1)(x-2), \dots, x(x-1)(x-2)\dots(x-n+1) \}$  is also a basis.

The set of all polynomials in  $x$  is a vector space which is infinite-dimensional.

A basis is  $\{ 1, x, x^2, x^3, x^4, \dots \}$

Examples of polynomials:  $5-3x+2x^2, 1-x^3+3x^7+11x^8, \dots$

Not polynomials:  $\sin x, \sqrt{1+x}, x - \frac{x^3}{6} + \frac{x^5}{120} - \frac{x^7}{5040} + \dots$

The set of all functions  $\mathbb{R} \rightarrow \mathbb{R}$ .

As a subspace of this, the continuous functions  $\mathbb{R} \rightarrow \mathbb{R}$ .

An even smaller subspace: differentiable functions  $\mathbb{R} \rightarrow \mathbb{R}$ .

Even smaller: the space of "smooth functions"  $V = \{ f: \mathbb{R} \rightarrow \mathbb{R} : f^{(n)} \text{ exists for all } n \geq 0 \}$

A linear transformation  $T: V \rightarrow V$  is defined by  $T = D^2 + I$  ( $D = \frac{d}{dx}$ ) i.e.  $Tf = f'' + f$ .

The rank of  $T$  is infinite dimensional.  $T$  is not one-to-one.

A basis for  $\text{Nul } T = \{ f: Tf = 0 \}$  is  $\{ \sin x, \cos x \}$ .

$Tf = 0$  iff  $f(x) = a \sin x + b \cos x$  for some  $a, b \in \mathbb{R}$ .

$D: V \rightarrow V$  has  $\text{Nul } D = \{ \text{constant functions} \}$  having basis  $\{ 1 \}$ ;  $\text{Nul } D$  is one-dimensional.

$D$  has eigenvectors! eg.  $D e^{3x} = 3e^{3x}$ . For every  $\lambda \in \mathbb{R}$ , the set of eigenvectors having eigenvalue  $\lambda$  is one-dimensional with basis  $\{ e^{\lambda x} \}$ .

# Fibonacci Numbers

0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, ...

Recursive formula  $F_n = \begin{cases} 0, & \text{if } n=0 \\ 1, & \text{if } n=1 \\ F_{n-1} + F_{n-2}, & \text{if } n \geq 2 \end{cases}$

$F_0 = 0$   
 $F_1 = 1$   
 $F_2 = 1$   
 $F_3 = 2$   
 $F_4 = 3$  etc.  
 $F_5 = 8$

Consider  $\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 2 \end{bmatrix}, \begin{bmatrix} 5 \\ 3 \end{bmatrix}, \dots$   
 $v_0 \quad v_1 \quad v_2 \quad v_3 \quad v_4$

So  $v_n = \begin{bmatrix} F_{n+1} \\ F_n \end{bmatrix}$  so  $A = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$  defines a map  $v_n \mapsto Av_n = v_{n+1}$  i.e.  $A \begin{bmatrix} F_{n+1} \\ F_n \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} F_{n+1} \\ F_n \end{bmatrix} = \begin{bmatrix} F_{n+2} \\ F_{n+1} \end{bmatrix} = v_{n+1}$ .

Starting with  $v_0 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ , we get  $v_1 = Av_0, v_2 = Av_1 = A^2v_0, \dots, v_n = \begin{bmatrix} F_{n+1} \\ F_n \end{bmatrix} = A^n \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \text{first column of } A^n$ .

$A^2 = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}, A^3 = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 2 & 1 \end{bmatrix}, A^4 = \begin{bmatrix} 3 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 5 & 3 \\ 3 & 2 \end{bmatrix}, \dots$

To find an explicit formula for  $A^n$  (and thereby  $F_n$ ), diagonalize  $A$ .

Characteristic polynomial of  $A$ :

$\det(A - xI) = \det \left( \begin{bmatrix} 1-x & 1 \\ 1 & -x \end{bmatrix} \right) = \begin{vmatrix} 1-x & 1 \\ 1 & -x \end{vmatrix} = (1-x)(-x) - 1 = x^2 - x - 1 = (x-\alpha)(x-\beta)$  where  $\alpha = \frac{1+\sqrt{5}}{2}$ ,  $\beta = \frac{1-\sqrt{5}}{2}$   
 golden ratio  $\approx 1.618\dots$   $-0.618\dots$

Eigenvector for  $\alpha$ : solution of  $Av = \alpha v$  i.e.  $(A - \alpha I)v = 0$

$\begin{bmatrix} 1-\alpha & 1 \\ 1 & -\alpha \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = 0$ . A nonzero solution is  $\begin{bmatrix} \alpha \\ 1 \end{bmatrix}$ . Check:  $\begin{bmatrix} 1-\alpha & 1 \\ 1 & -\alpha \end{bmatrix} \begin{bmatrix} \alpha \\ 1 \end{bmatrix} = \begin{bmatrix} (1-\alpha)\alpha + 1 \\ \alpha - \alpha \end{bmatrix} = \begin{bmatrix} 1 + \alpha - \alpha^2 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \approx 1.618\dots$

Eigenvector for  $\beta$ :  $Av = \beta v$  i.e.  $(A - \beta I)v = 0$ . Take  $\begin{bmatrix} \beta \\ 1 \end{bmatrix}$ .

$B = \begin{bmatrix} \alpha & \beta \\ 1 & 1 \end{bmatrix}$  has the eigenvectors as its columns.  $AB = \begin{bmatrix} \alpha & \beta \\ 1 & 1 \end{bmatrix} A \begin{bmatrix} \alpha \\ 1 \end{bmatrix} \begin{bmatrix} \beta \\ 1 \end{bmatrix} = \begin{bmatrix} \alpha^2 & \beta^2 \\ \alpha & \beta \end{bmatrix} = \begin{bmatrix} \alpha & \beta \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \alpha & 0 \\ 0 & \beta \end{bmatrix} = BD, D = \begin{bmatrix} \alpha & 0 \\ 0 & \beta \end{bmatrix}$ .

Diagonalizing  $A$  gives  $D = \begin{bmatrix} \alpha & 0 \\ 0 & \beta \end{bmatrix}$ .  $ABB^{-1} = BDB^{-1}$  i.e.  $A = BDB^{-1}$   $D^n = \begin{bmatrix} \alpha^n & 0 \\ 0 & \beta^n \end{bmatrix} = \begin{bmatrix} \alpha^n & 0 \\ 0 & \beta^n \end{bmatrix}$

$A^n = \underbrace{(BDB^{-1})(BDB^{-1})\dots(BDB^{-1})}_{n \text{ times}} = BD^nB^{-1}$

$B = \begin{bmatrix} \alpha & \beta \\ 1 & 1 \end{bmatrix}$   $B^{-1} = \frac{1}{\sqrt{5}} \begin{bmatrix} 1 & -\beta \\ 1 & \alpha \end{bmatrix}$   
 $\det B = \alpha - \beta = \sqrt{5}$

$$A^n = BDB^{-1} = \begin{bmatrix} \alpha & \beta \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \alpha^n & 0 \\ 0 & \beta^n \end{bmatrix} \begin{bmatrix} 1 & -\beta \\ 1 & \alpha \end{bmatrix}$$

$$= \frac{1}{\sqrt{5}} \begin{bmatrix} \alpha^{n+1} - \beta^{n+1} & \alpha\beta^n - \beta\alpha^{n+1} \\ \alpha\beta^n - \beta\alpha^{n+1} & \alpha\beta^n - \beta\alpha^{n+1} \end{bmatrix}$$

$\alpha^n - \beta^n$   
 $\alpha^{n+1} - \beta^{n+1}$

$$v_n = A^n \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} F_{n+1} \\ F_n \end{bmatrix} = \frac{1}{\sqrt{5}} \begin{bmatrix} \alpha^{n+1} - \beta^{n+1} \\ \alpha^n - \beta^n \end{bmatrix}$$

so

$$F_n = \frac{\alpha^n - \beta^n}{\sqrt{5}} = \frac{\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n}{\sqrt{5}}$$

(faster than power law  $n^k$ )

$$\alpha\beta = \left(\frac{1+\sqrt{5}}{2}\right)\left(\frac{1-\sqrt{5}}{2}\right) = \frac{1-5}{4} = \frac{-4}{4} = -1$$

$$\alpha\beta = -1$$

$$\alpha + \beta = 1$$

$$\alpha - \beta = \sqrt{5}$$

eg.  $F_0 = \frac{\alpha^0 - \beta^0}{\sqrt{5}} = \frac{1-1}{\sqrt{5}} = 0$

$$F_1 = \frac{\alpha^1 - \beta^1}{\sqrt{5}} = \frac{\sqrt{5}}{\sqrt{5}} = 1$$

$$F_2 = \frac{\alpha^2 - \beta^2}{\sqrt{5}} = \frac{(\alpha+1) - (\beta+1)}{\sqrt{5}} = \frac{\alpha - \beta}{\sqrt{5}} = 1$$

$$F_3 = 2 \text{ etc.}$$

$$F_{30} = \frac{\alpha^{30} - \beta^{30}}{\sqrt{5}} = 832040$$