


Analysis I (Math 3205)

Fall 2020

Book 2

{ Lebesgue integrable } \supseteq { Riemann integrable functions } \supseteq { continuous functions } \supseteq { differentiable functions } \supseteq { continuously differentiable functions } $\supseteq \dots$

\Downarrow
 g Dirichlet function H Heaviside function
 $H(x) = \begin{cases} 1 & \text{if } x \geq 0, \\ 0 & \text{if } x < 0 \end{cases}$



$$\int_{-2}^3 H(x) dx = 3$$

$S = \left\{ \frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \frac{5}{6}, \dots \right\}$ is bounded : 0 is a lower bound, 1 is an upper bound.
 $\frac{1}{2}$ is the greatest lower bound.



Every $m \leq \frac{1}{2}$ is a lower bound for S , meaning $s \geq m$ for all $s \in S$.


($0 \leq \frac{1}{2}$ is a lower bound) and it is the greatest lower bound.

1 is the least upper bound of S .

Fact: $|[0,1]| = |\mathbb{R}^3|$

Basic idea of the proof: $|(0,1)| = |(0,1)^3|$

$$(0,1)^3 = (0,1) \times (0,1) \times (0,1) = \{(x,y,z) : 0 < x, y, z < 1\}$$



Bijection: $a \mapsto (a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9, a_{10}, \dots)$

$$0 < a < 1$$

$$0.14159265358\dots \mapsto (0.1565\dots, 0.4958\dots, 0.1239\dots)$$

$$a = a_1 \cdot 10^0 + a_2 \cdot 10^{-1} + a_3 \cdot 10^{-2} + a_4 \cdot 10^{-3} + a_5 \cdot 10^{-4} + \dots$$

$$= a_1 \cdot 10^0 + a_2 \cdot 10^{-1} + a_3 \cdot 10^{-2} + a_4 \cdot 10^{-3} + a_5 \cdot 10^{-4} + \dots, \quad a_i \in \{0, 1, 2, \dots, 9\}$$

$$|(0,1)| = |\mathbb{R}|$$

see video on Cardinality

$|(0,1)| = |(0,1)^3| = |\mathbb{R}^3|$ and this bijection can be given constructively ie. by an explicit formula (in particular this is a theorem in ZF, not requiring the Axiom of Choice)




$$|[0,1]| = |[0,1]^2|$$



There is a bijection $[0,1] \rightarrow [0,1]^2$ but no continuous bijection.


However there is a continuous Surjection

This gives a "space-filling curve": (map that is onto) it goes through every point of the square.



etc.


How do you cut a hole in an $8\frac{1}{2} \times 11$ " sheet of paper that you can walk through?



→
fold in
half



→
open



Fact: There is a set of open intervals in \mathbb{R} of total length less than 1 which covers all the rational numbers.

Since \mathbb{Q} is countable, $\mathbb{Q} = \{q_1, q_2, q_3, q_4, q_5, \dots\}$. Then


$$\mathbb{Q} \subseteq \bigcup_{n=1}^{\infty} \left(q_n - \frac{1}{2^{n+1}}, q_n + \frac{1}{2^{n+1}}\right) = (q_1 - \frac{1}{4}, q_1 + \frac{1}{4}) \cup (q_2 - \frac{1}{8}, q_2 + \frac{1}{8}) \cup (q_3 - \frac{1}{16}, q_3 + \frac{1}{16}) \cup (q_4 - \frac{1}{32}, q_4 + \frac{1}{32}) \cup \dots$$

$$\text{Total length} < \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots = 1$$

Once again the set of intervals can be given constructively ie. explicitly with no need for the Axiom of Choice.

What is a (Riemann) integral? i.e. the integral, as defined in Calculus I-II?

Suppose $f: [a,b] \rightarrow \mathbb{R}$. We want to define $\int_a^b f(x) dx$. We start with lower and upper bounds for the integral (these being upper and lower Riemann sums). We then take $\sup \{\text{lower Riemann sums}\}$ and $\inf \{\text{upper Riemann sums}\}$.



First subdivide $[a,b]$ at points $a=x_0 \leq x_1 \leq x_2 \leq \dots \leq x_n=b$

giving n subintervals $[x_{i-1}, x_i]$, $i=1, 2, \dots, n$

Assuming f is bounded, $m_i \leq f(x) \leq M_i$ on $[x_{i-1}, x_i]$

$$\begin{aligned} M_i &= \sup \{f(x) : x_{i-1} \leq x \leq x_i\}, \quad m_i = \inf \{f(x) : x_{i-1} \leq x \leq x_i\} \\ &= \sup_{[x_{i-1}, x_i]} f \\ &= \inf_{[x_{i-1}, x_i]} f \end{aligned}$$

The Riemann Sums corresponding to the partition $a = x_0 \leq x_1 \leq x_2 \leq \dots \leq x_{n-1} \leq x_n = b$ of $[a, b]$ are :

$$\text{Upper sum } \sum_{i=1}^n \underbrace{(x_i - x_{i-1})}_{\text{base}} \underbrace{m_i}_{\text{height}} = (x_1 - x_0)M_1 + (x_2 - x_1)M_2 + \dots + (x_n - x_{n-1})M_n$$

$$\text{Lower sum } \sum_{i=1}^n (x_i - x_{i-1}) m_i$$

$$\text{We should have lower sum } \leq \int_a^b f(x) dx \leq \sum_{i=1}^n (x_i - x_{i-1}) m_i$$

We can't just let $n \rightarrow \infty$. By the Least Upper Bound Property, $\sup \{\text{lower bounds}\}$ exists and $\inf \{\text{upper bounds}\}$ exists. And

$$\sup \{\text{lower bounds}\} = \inf \{\text{upper bounds}\}$$

If these two values agree, this gives a definite value for $\int_a^b f(x) dx$.

For lots of functions (eg. the heaviside function and for all continuous functions), this works. For Dirichlet's function, the Riemann integral $\int_0^1 g(x) dx$ is undefined.

Why? For Dirichlet's function $g(x) = \begin{cases} 1, & \text{if } x \in \mathbb{Q}; \\ 0, & \text{if } x \notin \mathbb{Q} \end{cases}$

$$m_i = 0, \quad M_i = 1 \quad \text{for each } i = 1, 2, \dots, n$$


Upper sums: $\sum_{i=1}^n (x_i - x_{i-1}) \underbrace{M_i}_1 = (x_1 - x_0) + (x_2 - x_1) + (x_3 - x_2) + \dots + (x_m - x_{m-1}) + (x_n - x_{m-1})$
 $= x_n - x_0 = b - a = 1 - 0 = 1$

Lower sums: $\sum_{i=1}^m (x_i - x_{i-1}) \underbrace{m_i}_0 = 0 + 0 + \dots + 0 = 0$.

For C the Cantor Set define

$$u(x) = \begin{cases} 1, & \text{if } x \in C \\ 0, & \text{if } x \notin C \end{cases}$$

What is $\int_0^1 u(x) dx$?
 lower sums are 0.



For $0 < \frac{1}{3} < \frac{2}{3} < 1$ the Riemann Sum is $\frac{1}{3} \cdot 1 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = \frac{2}{3}$.

For $0 < \frac{1}{9} < \frac{2}{9} < \frac{1}{3} < \frac{2}{3} < \frac{7}{9} < \frac{8}{9} < 1$ the Riemann Sum is

$$\frac{1}{9} \cdot 1 + \frac{1}{9} \cdot 0 + \frac{1}{9} \cdot 1 + \frac{1}{3} \cdot 0 + \frac{1}{9} \cdot 1 + \frac{1}{9} \cdot 0 + \frac{1}{9} \cdot 1 = \frac{4}{9}$$

For $0 < \frac{1}{27} < \frac{2}{27} < \dots < \frac{26}{27} < 1$ the corresponding Riemann Sum is $\frac{8}{27}$.

Each new upper Riemann Sum is $\frac{2}{3}$ of the previous one if $\left(\frac{2}{3}\right)^n$.
 imp {upper Sums} = 0.

$$\text{For the function } u(x), \quad \underbrace{\sup \{ \text{lower sums} \}}_0 = \int_0^1 u(x) dx \leq \underbrace{\inf \{ \text{upper sums} \}}_0$$

$$\text{so } \int_0^1 u(x) dx = 0.$$

Note: $u(x)$ has infinitely many discontinuities but it is not discontinuous everywhere.

$u(x)$ is continuous on a set of open intervals inside $[0, 1]$ of total length 1.

The total length of the Cantor set C (where $\kappa=1$) is 0.

However C is uncountable; $|C| = |\mathbb{R}|$. Why?

Every $a \in [0, 1]$ has a ternary expansion

$$a = 0.a_1 a_2 a_3 a_4 a_5 \dots \quad (a_i \in \{0, 1, 2\})$$


$$= \frac{a_1}{3} + \frac{a_2}{3^2} + \frac{a_3}{3^3} + \frac{a_4}{3^4} + \frac{a_5}{3^5} + \dots$$

The points in C are those with $a_i \in \{0, 2\}$ only.

$|C| = |\mathbb{R}| (= |[0, 1]|)$. A bijection $C \rightarrow [0, 1]$

$$0.20021102002 \dots \xrightarrow{\text{(base 3) ternary}} 0.10011101001 \dots \xrightarrow{\text{(base 2) binary}}$$

If two functions f and g agree except at a single point, then $\int_a^b f(x) dx = \int_a^b g(x) dx$



The same holds for changing a function at any finite number of points.