

Algebra I

Group Theory

Book 2

Transpositions (ij) are odd permutations.

$$(123456789) = (19)(18)(17)(16)(15)(14)(13)(12)$$

A k -cycle is a product of $k-1$ transpositions.

If k is even, this is odd; and vice versa.

A cycle of odd length is an even permutation;
 even odd

If α is a product of an even number of transpositions, then α is an even permutation.
 odd odd

Permutations in S_5 :


Even	
$()$	1
(ijk)	20
$(ijklm)$	24
$(ij)(kl)$	15
	<hr/>
	60

Odd	
(ij)	10
$(ijkl)$	30
$(ijk)(lm)$	20
	<hr/>
	60

$$|S_5| = 120$$

$$A_5 = \{ \text{even permutations in } S_5 \}$$

$$|A_5| = 60$$



An even permutation of the coordinate axis in \mathbb{R}^n is an orientation-preserving transformation.

An odd permutation of the coordinate axis in \mathbb{R}^n is an orientation-reversing transformation.

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a linear transformation then

$$\det T \begin{cases} = 0 & \text{if } T \text{ is not invertible} \\ > 0 & \text{preserves orientation} \\ < 0 & \text{reverses} \end{cases}$$

A permutation $\alpha \in S_n$ can be expressed as a product of transpositions.

If α is a product of an even number of transpositions, then α is even.

In S_3 :

$(13)(12)(13)(23)(23)(12)(23) = (123)$ says (123) is an even permutation.

$S_3 \cong \langle \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ -1 & -1 \end{bmatrix} \rangle \cong$ dihedral group of order 6
(symmetry group of an equilateral triangle)

Groups of order 2

$S_2 \cong \{0, 1\} \pmod 2$ under addition $\cong \langle -1 \rangle$ under multiplication

n	no. of groups of order n up to isomorphism
1	1
2	1
3	1
4	2
5	1
6	2
7	1
8	5

o	(1)	(12)	+	0	1	.	1	-1
(1)	(1)	(12)	0	0	1	1	1	-1
(12)	(12)	(1)	1	1	0	-1	-1	1



has a cyclic symmetry group of order 4



has an abelian symmetry group of order 4 which is not cyclic (the Klein four-group)

Cayley tables of groups of order 2 all "look the same"

Theorem Any two groups of prime order are isomorphic; they are cyclic of order p.

Eg. $\mathbb{Z}/3\mathbb{Z} = \{0, 1, 2\}$ (under addition mod 3) is isomorphic to $A_3 = \langle (123) \rangle = \{(), (123), (132)\}$ and $\{1, \omega, \omega^2\}$ under multiplication, $\omega = \frac{-1+i\sqrt{3}}{2} = e^{2\pi i/3}$


\oplus	0	1	2
0	0	1	2
1	1	2	0
2	2	0	1

\circ	$()$	(123)	(132)
$()$	$()$	(123)	(132)
(123)	(123)	(132)	$()$
(132)	(132)	$()$	(123)

\cdot	1	ω	ω^2
1	1	ω	ω^2
ω	ω	ω^2	1
ω^2	ω^2	1	ω



We say two groups G, H are isomorphic ($G \cong H$) if there exists a bijection $\phi: G \rightarrow H$ such that $\phi(xy) = \phi(x)\phi(y)$



operation in G operation in H

An isomorphism $\phi: \mathbb{Z}/3\mathbb{Z} \rightarrow A_3$ is a bijection satisfying $\phi(x+y) = \phi(x)\phi(y)$

An isomorphism $\phi: \mathbb{R} \xrightarrow{\text{under addition}} (0, \infty) \xrightarrow{\text{under multiplication}}$ is defined by $\phi(x) = e^x$
 $e^{x+y} = e^x \cdot e^y$

(subgroup of $\mathbb{R}^* = (-\infty, 0) \cup (0, \infty)$)
 $\ln = \phi^{-1}: (0, \infty) \rightarrow \mathbb{R}$

$\mathbb{R} \not\cong \mathbb{R}^*$
 since \mathbb{R} (reals under addition) has only one element of finite order whereas \mathbb{R}^* has two elements of finite order: ± 1 .

+	0	1	2
0	0	1	2
1	1	2	0
2	2	0	1

is isomorphic to

*	a	b	c
a	b	c	a
b	c	a	b
c	a	b	c

$$\begin{aligned} \phi(0) &= c \\ \phi(1) &= a \\ \phi(2) &= b \end{aligned} \quad * \begin{array}{c|ccc} & c & a & b \\ \hline c & c & a & b \\ a & a & b & c \\ b & b & c & a \end{array}$$

or

$$\begin{aligned} \phi(0) &= c \\ \phi(1) &= b \\ \phi(2) &= a \end{aligned} \quad * \begin{array}{c|ccc} & c & b & a \\ \hline c & c & b & a \\ b & b & a & c \\ a & a & c & b \end{array}$$

$\mathbb{Z}/3\mathbb{Z}$

Every group of order 1 is isomorphic to $\mathbb{Z}/1\mathbb{Z}$
 2 $\mathbb{Z}/2\mathbb{Z}$

+	0	1
0	0	1
1	1	0

(trivial group $\{1\}$)

	c
a	ac
b	bc

If $ac=bc$ then multiply both sides by c^{-1} on the right
 to get $(ac)c^{-1} = (bc)c^{-1}$
 $a(cc^{-1}) = b(cc^{-1})$
 $a1 = b1$
 $a = b$

	e	a	b
e	e	a	b
a	a	b	e
b	b	e	a

Every group of order 3 is cyclic (isomorphic to $\mathbb{Z}/3\mathbb{Z}$ under addition).

	e	a	b	c
e	e	a	b	c
a	a	e	c	b
b	b	c	e	a
c	c	b	a	e

Klein four-group

	e	a	b	c
e	e	a	b	c
a	a	b	c	e
b	b	c	e	a
c	c	e	a	b

Cyclic group of order 4

Two cases: either all ^{non-identity} elements of G have order 2, or G has an element not of order 2.

Theorem: There are exactly two groups of order 4 up to isomorphism: the Klein four-group and the cyclic group of order 4.

	e	a	b	c	d
e	e	a	b	c	d
a	a	b	c	d	e
b	b	c	d	e	a
c	c	d	e	a	b
d	d	e	a	b	c

cyclic group of order 5

$$\langle a \rangle = \{e, a, a^2, a^3, a^4\}$$

$\begin{matrix} & \uparrow & \uparrow & \uparrow \\ & b & c & d \end{matrix}$

	e	a	b	c	d
e	e	a	b	c	d
a	a	e	c	d	b
b	b	c	d	a	e
c	c	d	e	b	a
d	d	b	a	e	c

c is a left inverse for b ($cb=e$) but not a right inverse for b ($bc=a$).

is not a group!

It is a quasigroup, in fact since it has an identity e , it is a loop (its Cayley table is a Latin square: each row/column is a permutation of e, a, b, c, d).

This loop is not associative eg. $(ca)d = dd = c$
 $c(ad) = cb = e$

Theorem If every ^{non-identity} element of a group G has order 2, then G is abelian.

Proof (Note: $x^2=e$ = identity for every $x \in G$.)

Let $x, y \in G$. Then $(xy)^2 = xyxy = e$ so

$$yx = \underbrace{x(xyxy)}_{x^2=e} \underbrace{y}_{y^2=e} = xey = xy. \quad \square$$

\curvearrowright In such groups, $x^{-1} = x$ for all $x \in G$.

Shoe-Sock Theorem

In every group G , with identity 1 , for $x, y \in G$ we have $(xy)^{-1} = y^{-1}x^{-1}$.

Proof $(y^{-1}x^{-1})(xy) = y^{-1}1y = 1$ and $(xy)(y^{-1}x^{-1}) = 1$. \square

Warning: $(xy)^{-1} \neq x^{-1}y^{-1}$ in general.

	e	a	b	c
e	e	a	b	c
a	a	e	c	b
b	b	c	e	a
c	c	b	a	e

Klein four-group

	e	a	b	c
e	e	a	b	c
a	a	b	c	e
b	b	c	e	a
c	c	e	a	b

Cyclic group of order 4

Write the rows of the Cayley table as permutations of $\overset{1}{e}, \overset{2}{a}, \overset{3}{b}, \overset{4}{c}$:
 $\{(1), (12)(34), (13)(24), (14)(23)\}$ is a Klein four group as a subgroup of S_4 .

Gives $\{(1), (1234), (13)(24), (1432)\}$ as a subgroup of S_4 .

Theorem (Cayley Representation Theorem)
 Every finite group G is isomorphic to a subgroup of S_n where $n = |G|$.

By the way, every finite group G is also isomorphic to a group of matrices under multiplication.

Theorem If G is a finite group of order n , then every element $g \in G$ has order dividing n .
(If $g \in G$ then $|g| \mid n$.)

Eg. S_4 has elements of order 1, 2, 3, 4. These orders of elements divide $|S_4| = 24$.

S_5 has elements of order 1, 2, 3, 4, 5, 6 (divisors of $|S_5| = 120$).

Proof In the general case this follows from a later theorem, Lagrange's Theorem. Here let's prove the theorem in the special case that G is abelian. (We have already proved the result for cyclic groups.)

Consider the product of all the group elements $\pi = g_1 g_2 \dots g_n$ where $G = \{g_1, g_2, \dots, g_n\}$, $g_1 = 1$.

Note: since G is abelian, π is well-defined; it doesn't depend on what order we list the elements $g_1, \dots, g_n \in G$. Pick $a \in G$. (So $a \in \{g_1, \dots, g_n\}$.) The elements ag_1, ag_2, \dots, ag_n

are again all the elements of G so

$$(ag_1)(ag_2)(ag_3) \dots (ag_n) = \pi = a^n g_1 g_2 \dots g_n = a^n \pi$$


so $a^n = 1$ and $k = |a|$ must divide n . \square

Lagrange's Theorem If G is any finite group of order n , and $H \leq G$ (i.e. H is a subgroup of G) then $|H| \mid n$.

This generalizes the previous statement: if $g \in G$ then by Lagrange's Theorem, $| \langle g \rangle | = |g| \mid |G|$.

Eg. $|A_4| = \frac{1}{2} |S_4| = 12$, $A_4 = \{ (1), (123), (124), (132), (134), (142), (143), (234), (243), (12)(34), (13)(24), (14)(23) \}$.

The symmetry group of a regular tetrahedron



is isomorphic to S_4 .

The rotational symmetry group of the regular tetrahedron (the direct isometry group, consisting of those symmetries that preserve orientation) is isomorphic to A_4 .

$$A_4 = \{(), (123), (124), (132), (134), (142), (143), (234), (243), (12)(34), (13)(24), (14)(23)\}.$$

Subgroups of A_4 have order 1, 2, 3, 4.

Elements of A_4 have order 1, 2, 3.

Divisors of $|A_4| = 12$ are 1, 2, 3, 4, 6, 12.

$$\langle (243), (12)(34) \rangle = \{(), (243), (12)(34), (234), (142), (124), \dots\} = A_4.$$

$$(243)(12)(34) = (142)$$

$\{(), (12)(34), (13)(24), (14)(23)\}$ is the Klein four-group, a subgroup of A_4 .

Question: How many subgroups of \mathbb{Z} are there containing 4? (Note: \mathbb{Z} is an additive group.)

Answer: There are three subgroups of \mathbb{Z} containing 4, namely \mathbb{Z} , $2\mathbb{Z}$, $4\mathbb{Z}$.

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, 4, 5, \dots\}$$

$$2\mathbb{Z} = \{\dots, -6, -4, -2, 0, 2, 4, 6, 8, \dots\}$$

$$4\mathbb{Z} = \{\dots, -8, -4, 0, 4, 8, 12, \dots\}$$

$$-4\mathbb{Z} = \{\dots, -8, -4, 0, 4, 8, 12, \dots\}$$

\mathbb{Z} has infinitely subgroups: one finite subgroup $\{0\}$ and all the other subgroups are infinite.

There are infinite subgroups of \mathbb{Z} containing 4 but not infinitely many subgroups of \mathbb{Z} containing 4.

Note: For every cyclic group G , all subgroups of G are cyclic; they are generated by powers of the generator of G .

Ex. $G = \langle g \rangle$ where $|g| = \infty$ i.e. $|G| = |\langle g \rangle| = |g| = \infty$.

$= \{ \dots, g^{-3}, g^{-2}, g^{-1}, 1, g, g^2, g^3, \dots \}$ with no repeats.

1 is the identity


$$g^i g^j = g^{(i+j)} = g^j g^i$$

How many subgroups of $G = \langle g \rangle$ contain g^4 ? Three: $\langle g \rangle, \langle g^2 \rangle, \langle g^4 \rangle$.

$$G = \{ \dots, g^{-3}, g^{-2}, g^{-1}, 1, g, g^2, g^3, g^4, \dots \}$$

$$\langle g^2 \rangle = \{ \dots, g^6, g^4, g^2, 1, g^2, g^4, g^6, \dots \}$$

$$\langle g^4 \rangle = \{ \dots, g^8, g^4, 1, g^4, g^8, g^{12}, \dots \}$$



$$\langle g^6, g^{10} \rangle \leq \langle g^2 \rangle$$

$$\langle g^2 \rangle \leq \langle g^6, g^{10} \rangle$$

Since $g^2 = (g^6)^2 (g^{10})^{-1}$

So $\langle g^2 \rangle = \langle g^6, g^{10} \rangle$

$G \cong \mathbb{Z}$
 multiplicative cyclic group \cong additive cyclic group

$\phi: \mathbb{Z} \rightarrow G$ is an isomorphism
 $\phi(i) = g^i$

Theorem If G is a group of even order, then G has an element of order 2 (i.e. at least one element of order 2). Note: G is not necessarily abelian.

Proof Pair up each group element with its inverse giving pairs $\{g, g^{-1}\}$ for $g \in G$. Note that $g = g^{-1}$ iff g has order 1 or 2. ($g = g^{-1} \iff g^2 = 1 \iff |g|$ divides 2). So G is partitioned into subsets $\{g, g^{-1}\}$ having size 1 or 2. If G has no elements of order 2 then we have partitioned a set G of even cardinality into one subset $\{1\}$ of size 1, and a collection of pairs $\{g, g^{-1}\}$ of size 2, a contradiction. \square

what we actually showed is that in a group of even order, the number of elements of order 2 is odd. (In a group of odd order, there are no elements of order 2 although we haven't proved this yet except in the abelian case.)

Ex. Direct Products: Given groups G, H (say, multiplicative) we form the direct product of G and H as $G \times H = \{(g, h) : g \in G, h \in H\}$ (the cartesian product of the sets G and H) which becomes a group under coordinatewise multiplication i.e.

$$(g, h)(g', h') = (gg', hh')$$

and coordinatewise inverses i.e. $(g, h)^{-1} = (g^{-1}, h^{-1})$

and the coordinatewise identity $1 \in G \times H$ is $1 = 1_{G \times H} = (1_G, 1_H)$. or $e_{G \times H} = (e_G, e_H)$.

Ex. $\mathbb{Z}/2\mathbb{Z} = \{0, 1\}$ under addition mod 2

$$\begin{array}{c|c} + & \begin{array}{c} 0 \\ 1 \end{array} \\ \hline \begin{array}{c} 0 \\ 1 \end{array} & \begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array} \end{array}$$

$$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} = \{(x, y) : x, y \in \mathbb{Z}/2\mathbb{Z}\} = \{(0, 0), (0, 1), (1, 0), (1, 1)\}$$

$$(x, y) + (x', y') = (x+x', y+y'). \quad \text{The identity } 0 = (0, 0).$$

This is the Klein four-group since it has 3 elements of order 2.

Note: Many books write \mathbb{Z}_2 in place of $\mathbb{Z}/2\mathbb{Z}$ or \mathbb{Z}_2

If $|G|=m$ and $|H|=n$ then $|G \times H| = mn$.

If G and H are abelian then so is $G \times H$.

In fact, the converse holds: G and H are both abelian, iff $G \times H$ is abelian.

$$G \times H \cong H \times G$$

$$\phi: G \times H \rightarrow H \times G$$

$$\phi(g, h) = (h, g) \text{ is an isomorphism.}$$

$G \times H$ has a subgroup $G \times \{1_H\} = \{(g, 1_H) : g \in G\} \cong G$

An isomorphism $G \times \{1_H\} \rightarrow G$ is given by $(g, 1_H) \mapsto g$.

Like wise, $G \times H$ has a subgroup $\{1_G\} \times H \cong H$

$$(g, 1_H)(1_G, h) = (g, h) = (1_G, h)(g, 1_H)$$

$$\begin{array}{ccc} \underbrace{G \times \{1_H\}} & & \underbrace{\{1_G\} \times H} \\ \uparrow & & \uparrow \\ G & & H \end{array}$$

Eg. $\mathbb{R}^* = (-\infty, 0) \cup (0, \infty) \cong \underbrace{\mathbb{R}}_{\text{additive group}} \times \underbrace{\mathbb{Z}/2\mathbb{Z}}_{\text{additive}}$
multiplicative group

An isomorphism $\phi: \mathbb{R}^* \rightarrow \mathbb{R} \times \mathbb{Z}/2\mathbb{Z}$ is $\phi(a) = \begin{cases} (\ln|a|, 0) & \text{if } a > 0 \\ (\ln|a|, 1) & \text{if } a < 0 \end{cases}$

It's easy to see that ϕ is one-to-one and onto.

We show that $\phi(ab) = \phi(a) + \phi(b)$ for all $a, b \in \mathbb{R}^*$.

We argue in four cases. If $a, b > 0$ then

$$\begin{aligned} \phi(ab) &= (\ln|ab|, 0) \quad \text{since } ab > 0 \\ &= (\ln|a| + \ln|b|, 0) = (\ln|a|, 0) + (\ln|b|, 0) = \phi(a) + \phi(b) \end{aligned}$$

If $a > 0 > b$ then $ab < 0$ so

$$\phi(ab) = (\ln|ab|, 1) = (\ln|a|, 0) + (\ln|b|, 1) = \phi(a) + \phi(b)$$

Similarly if $a < 0 < b$.

If $a, b < 0$ then $ab > 0$ so

$$\begin{aligned} \phi(ab) &= (\ln|ab|, 0) = (\ln|a|, 1) + (\ln|b|, 1) \\ &= \phi(a) + \phi(b) \end{aligned}$$

Every cyclic group is abelian.
 Not every abelian group is cyclic but every abelian group is a direct product of cyclic groups.
 eg. the Klein four-group is a direct product of two groups of order 2 i.e. $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$

There are five groups of order 8 up to isomorphism:

$\mathbb{Z}/8\mathbb{Z}$ (cyclic)

$$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} = \{(a,b) : a \in \mathbb{Z}/2\mathbb{Z}, b \in \mathbb{Z}/4\mathbb{Z}\}$$

$$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} = \{(a,b,c) : a,b,c \in \mathbb{Z}/2\mathbb{Z}\} \text{ under addition}$$

} three abelian groups of order 8

dihedral group of order 8 \cong symmetry group of square, D_4 (sometimes D_8)

quaternion group of order 8, Q or Q_8

$$Q = \{ \underbrace{1, -1}_{\text{order 2}}, \underbrace{i, -i, j, -j, k, -k}_{\text{order 4}} \} \quad \begin{array}{l} ij=k, \quad ji=-k, \quad i^2=j^2=k^2=-1 \\ jk=i, \quad kj=-i \\ ki=j, \quad ik=-j \end{array}$$

For any field F (eg. $\mathbb{R}, \mathbb{C}, \mathbb{Q}$) $GL_n(F) = \{ \text{invertible } n \times n \text{ matrices over } F \}$ i.e. having entries in F .

Also $F = \mathbb{F}_3 = \{0, 1, 2\}$ works with addition mod 3. $2+2=1=2 \times 2$
 $\frac{1}{2} = 2$

In $\mathbb{F}_7 = \{0, 1, 2, \dots, 6\}$, $\frac{1}{5} = 3$.

$\mathbb{F}_p = \{0, 1, 2, \dots, p-1\}$ is a field whenever p is prime.

$GL_2(\mathbb{F}_3) = \{ \text{invertible } 2 \times 2 \text{ matrices over } \mathbb{F}_3 \}$ is a group of order 48.

$GL_2(\mathbb{R}) = \{ \text{invertible } 2 \times 2 \text{ matrices over } \mathbb{R} \} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a,b,c,d \in \mathbb{R}, ad-bc \neq 0 \right\}$

$GL_n(F) = \{ \text{invertible } n \times n \text{ matrices over } F \} = \text{general linear group of degree } n \text{ over } F$
 also denoted $GL(n, F)$ in the textbook

$SL_n(F)$ is the special linear group of degree n over F ; $SL_n(F) \subseteq GL_n(F)$
 or $SL(n, F)$ $SL_n(F) = \{n \times n \text{ matrices over } F \text{ having determinant } 1\}$.

If $F = \mathbb{F}_p = \{0, 1, 2, \dots, p-1\}$ mod p (field of prime order p) then we can count elements in $GL_n(\mathbb{F}_p)$ or $SL_n(\mathbb{F}_p)$. (For 2×2 matrix over \mathbb{F}_3 , 33 matrices have $\det A = 0$, $\frac{24}{24}$ matrices have $\det A = 1$, $\frac{24}{24}$... $\det A = 2$).

$|GL_2(\mathbb{F}_3)| = 48$.

The number of 2×2 matrices over $\mathbb{F}_3 = \{0, 1, 2\}$ is 81. How many of them are invertible?

We count invertible matrices $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$, $a, b, c, d \in F = \mathbb{F}_3$ with linearly independent columns.

There are 8 choices for the first column $\begin{bmatrix} a \\ c \end{bmatrix} \neq \begin{bmatrix} 0 \\ 0 \end{bmatrix}$. $9 - 3 = 6$

Having chosen the first column $\begin{bmatrix} a \\ c \end{bmatrix}$, there are 6 choices for the second column $\begin{bmatrix} b \\ d \end{bmatrix}$ which are not a scalar multiple of the first column. So $|GL_2(\mathbb{F}_3)| = 8 \times 6 = 48$.

In fact, for $A \in GL_2(F)$, $F = \mathbb{F}_3$, there are 24 choices with determinant 1, and 24 choices with determinant $-1 = 2$.

$$|GL_n(\mathbb{F}_p)| = \underbrace{(p^n - 1)}_{\substack{\uparrow \\ \text{no. of choices} \\ \text{of first column}}} \underbrace{(p^n - p)}_{\substack{\uparrow \\ \text{no. of choices} \\ \text{of second column}}} \underbrace{(p^n - p^2)}_{\substack{\uparrow \\ \text{no. of choices of third column}}} \cdots \underbrace{(p^n - p^{n-1})}_{\substack{\uparrow \\ \text{no. of choices of} \\ \text{last column}}}$$

$|GL_2(\mathbb{F}_p)| = (p^2 - 1)(p^2 - p)$


For $A \in GL_n(\mathbb{F}_p)$, $\det A \in \{1, 2, \dots, p-1\}$ and there are equally many matrices with each possible nonzero determinant in $\{1, 2, \dots, p-1\}$ so

$|SL_n(\mathbb{F}_p)| = \frac{1}{p-1} |GL_n(\mathbb{F}_p)|$. We'll explain later.

For any group G , the center of G is $Z(G) = \{ \text{all elements in } G \text{ which commute with everything in } G \}$
 Zentrum \uparrow not Z = $\{ z \in G : zx = xz \text{ for all } x \in G \}$
 Zahlen

Ex. if G is the symmetry group of a square (a dihedral group of order 8) then $|Z(G)| = 2$
 and $Z(G)$ consists of the identity and the half-turn (180° rotation about the center).

If we represent G using permutations on the vertices 1, 2, 3, 4 then
 $G = \{ (), (1234), (13)(24), (1432), (12)(34), (14)(23), (13), (24) \}$
 then $Z(G) = \langle (13)(24) \rangle = \{ (), (13)(24) \}$.



Alternatively, G can be represented as a subgroup of $GL_2(\mathbb{R})$:

$$G = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 0 & -1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \right\}$$

$$Z(G) = \left\langle \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \right\rangle = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \right\}$$

In general, $Z(G) \leq G$ (a subgroup of G).
 $Z(G) = G$ iff G is abelian.

For many groups, $Z(G) = \{ 1 \}$ identity eg. $Z(S_3) = \{ () \}$.

Theorem If G is a group and $z \in G$, then $Z(G) \leq G$ (the center of G is a subgroup of G).
 $e = \text{identity of } G$

Proof Since $eg = g = ge$ for every $g \in G$, $e \in Z(G)$. If $z, z' \in Z(G)$ then

$$(zz')g = z(z'g) = z(gz') = (zg)z' = (gz)z' = g(zz')$$


so $zz' \in Z(G)$. Also if $z \in Z(G)$ then for every $g \in G$ we have $zg = gz$ so $z^{-1}g = z^{-1}(gz)z^{-1} = z^{-1}(zg)z^{-1} = gz^{-1}$

so $z^{-1} \in Z(G)$. \square

Another construction of subgroups: Suppose $G \leq S_n$. So G permutes $[n] = \{1, 2, \dots, n\}$.

The stabilizer of a point $x \in [n]$ is $\text{Stab}_G(x) = \{g \in G : g(x) = x\} \leq G$.

Eg.



The symmetry group of a regular pentagon is a group G which is dihedral of order 10 (sometimes denoted D_5 or D_{10}).

$$G = \{(), (12345), (13524), (14253), (15432), (12)(35), (13)(45), (14)(23), (15)(24), (25)(34)\}$$

5 rotations

5 reflections

$G \leq S_5$ permuting $[5] = \{1, 2, 3, 4, 5\}$, the five vertices.

$$\text{Stab}_G(3) = \{(), (15)(24)\}.$$

$$() (x) = x$$

If $g, h \in \text{Stab}_G(x)$ then

$$(gh)(x) = g(h(x)) = g(x) = x$$

If $g \in \text{Stab}_G(x)$ then $g(x) = x$ so

$$x = g^{-1}(g(x)) = g^{-1}(x) \quad \text{so} \quad g^{-1} \in \text{Stab}_G(x).$$

Elements of order 2 in a group are called involutions. If $|a|=|b|=2$ then $(ab)^2 = abab = a^2b^2 = 1 \cdot 1 = 1$ so $|ab|=1$ or 2. If $ab=1$ then $b=a$; otherwise $ab \neq 1$, $(ab)^2=1$ so ab is an involution so $\{1, a, b, ab\}$ is a Klein four-subgroup of G . Any two distinct involutions in G generate a Klein four-subgroup. $\langle a, b \rangle$ is an abelian group. $(12)(13) = (132)$ in S_3 .

How many involutions can a finite abelian group G have?

If G has k involutions then every involution lies in exactly $\frac{k-1}{2}$ Klein four-subgroups




How many Klein four-subgroups does G have altogether?

Count subgroups of the form $\langle a, b \rangle = \{1, a, b, ab\}$ where $a, b \in G$ are distinct involutions.

k choices for a
 $k-1$ choices for b

$\frac{k(k-1)}{6}$ is the number of Klein four-subgroups in G .



If $k=7$ then we have 7 involutions, 7 Klein four-subgroups, every involution is in 3 Klein four-groups, every Klein four-group has 3 involutions.

In a direct product of three groups of order two e.g. $\langle -1 \rangle \times \langle -1 \rangle \times \langle -1 \rangle$
 $= \{ (x, y, z) : x, y, z \in \langle -1 \rangle \}$

$$\langle -1 \rangle = \{1, -1\}$$

Certainly $k \equiv 1$ or $3 \pmod{6}$

In general if a, b are distinct involutions in a group G then what can they generate?

$$\langle a, b \rangle = \{1, a, b, ab, ba, aba, bab, abab, baba, \dots\} \text{ with possible duplicates.}$$


The symmetry group of an infinite string $\dots TTTT \dots$ is generated by two reflections a, b in vertical axes l, l' as shown

ab is a translation (shift) one step to the right
 ba is a translation one step to the left.

$\langle ab \rangle = \{\dots, baba, ba, 1, ab, abab, ababab, \dots\}$ is an infinite cyclic group, a subgroup of $\langle a, b \rangle$
 $\langle a, b \rangle$ itself is an infinite dihedral group.

The symmetry group of a square is a dihedral group $\langle R, R' \rangle$ generated by two reflections

$$\langle R, R' \rangle = \{I, R, R', RR', R'R, RR'R, R'R'R', RR'R'R, R'R'R'R\}$$



Comments on HW2:

Recall in class we used the product π of elements in a finite abelian group.

#5(a) Show that π has order ≤ 2 .

Proof If $G = \{g_1, g_2, \dots, g_n\}$ is abelian of order n then $\pi = g_1 g_2 \dots g_n = g_1^{-1} g_2^{-1} \dots g_n^{-1}$ so
 $\pi^2 = (\cancel{g_1} \cancel{g_2} \dots \cancel{g_n}) (\cancel{g_1^{-1}} \cancel{g_2^{-1}} \dots \cancel{g_n^{-1}}) = e$ (the identity element of G).

Eg. G is cyclic of order 4.

In multiplicative notation, $G = \langle g \rangle = \{1, g, g^2, g^3\}$ where $g^4 = 1$; $\pi = 1 \cdot g \cdot g^2 = g^3$ of order 2.

In additive notation, $G = \mathbb{Z}/4\mathbb{Z} = \{0, 1, 2, 3\} = \langle 1 \rangle$; $\pi = 0 + 1 + 2 + 3 = 2$ of order 2. 0 is the identity.

In S_4 , $G = \langle (1234) \rangle \leq S_4$, $\pi = () \circ (1234) \circ (13)(24) \circ (1432) = (13)(24)$ of order 2.

$\lim_{x \rightarrow \infty} \frac{\sin x}{x} = 0$. $\lim_{\varepsilon \rightarrow \infty} \frac{\sin \varepsilon}{\varepsilon} = 0$ is problematic in its unorthodox choice of variable $\varepsilon \rightarrow \infty$.

$\lim_{\varepsilon \rightarrow 0} \frac{\sin \varepsilon}{\varepsilon} = 1$ is natural.

$G = SL_2(\mathbb{F}_3) = \{2 \times 2 \text{ matrices of } \mathbb{F}_3 \text{ having determinant } 1\}$. $|G| = 24$.

Is $G \cong S_4$? G has only one involution whereas S_4 has 9 involutions. (An involution in any group element of order 2.)

If $G = SL_2(\mathbb{R})$ or $SL_2(\mathbb{C})$ then G has only one involution, $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} = -I$. $GL_2(\mathbb{R})$ has many involutions.

Does $GL_2(\mathbb{R})$ have an element of order 11? Yes; in fact

eg. $\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$ is a reflection in the y-axis.

$SL_2(\mathbb{R})$ does: $\begin{bmatrix} \cos \frac{2\pi}{11} & -\sin \frac{2\pi}{11} \\ \sin \frac{2\pi}{11} & \cos \frac{2\pi}{11} \end{bmatrix} \in SL_2(\mathbb{R})$

$A = \begin{bmatrix} \cos \theta & -\sin \theta \\ -\sin \theta & -\cos \theta \end{bmatrix} \in GL_2(\mathbb{R})$ has determinant -1 .

Why is $-I$ the only involution in $SL_2(\mathbb{R})$?

$$A^2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

A is a reflection

(Infinitely many involutions in $GL_2(\mathbb{R})$.)