

Combinatorics

Book 2

The graph $\Gamma \setminus v$ (formed by removing v and its edges from Γ) has one fewer vertex, so it can be properly colored using at most 6 colors. And since v has at most 5 neighbors in $\Gamma \setminus v$, there is a color left over which can be used to color vertex v . This gives a proper coloring of Γ using at most 6 colors (a contradiction...)

We will improve this to show that actually 5 colors suffice to properly color every planar graph.

$$\nexists \phi \chi \pi$$

Given a graph Γ , the chromatic number of Γ , denoted $\chi(\Gamma)$,

is the smallest number of colors we can use to properly color the vertices of Γ . A proper coloring of the vertices of Γ is a coloring of the vertices such that no edge has both endpoints of the same color.

A b c ... x x

↑

Greek "cli"

The theorem of Appel and Haken (1976) is that every planar graph Γ has $\chi(\Gamma) \leq 4$.

Note that $\chi(K_n) = n$. Here K_n is the complete graph of order n .


A graph Γ has $\chi(\Gamma) = 1$ iff it has vertices but no edges.

A graph Γ has $\chi(\Gamma) \leq 2$ iff Γ is bipartite iff Γ has no circuits of odd length.

Computing $\chi(\Gamma)$ is hard in general.

Theorem If Γ is a finite planar graph then $\chi(\Gamma) \leq 5$. Proof due to Heawood.

Proof If the theorem fails then there is a smallest counterexample Γ with n vertices (so Γ is planar and every planar graph of order $n-1$ has chromatic number ≤ 5 while $\chi(\Gamma) \geq 6$). We seek a contradiction. Γ has a vertex v of degree ≤ 5 . In fact $\deg v = 5$. (If $\deg v \leq 4$ then $\chi(\Gamma) \leq 5$, a contradiction.) Let Γ' be the graph obtained



from Γ by deleting v and its five edges,


so $\chi(\Gamma') \leq 5$. Say v_i has color i ($i=1, 2, \dots, 5$).

Consider the vertices $V_{13} \subset \{\text{vertices of } \Gamma\}$ having colors 1, 3 only. This graph is bipartite. I can assume v_1 is joined to v_3 in V_{13} (otherwise



$$\Gamma_{13} = \Gamma(V_{13})$$

in part of V_{13} , reverse colors 1, 3 so that v_3 gets color 1. Then we are free to color v using color 3 since its neighbors are color 1, 2, 4, 5).


Otherwise Γ_{13} has a path from v_1 to v_3 .




Similarly there is a path from v_2 to v_4 using only vertices of colors 2 and 4. Contradiction! \square



Given a graph Γ , a subgraph of Γ is formed by taking a subset of the edges of Γ together with all their vertices. An induced subgraph of Γ is formed by taking a subset of the vertices of Γ together with all their edges in Γ .



is a subgraph of Γ . (not an induced subgraph of Γ)



is an induced subgraph of Γ .

An induced subgraph of Γ is a subgraph of Γ , but not conversely.

A k -clique in Γ is a complete subgraph of Γ , i.e. a subset of the vertices, any two of which are joined.

In Γ above, $\{1, 2, 6\}$ is a clique (in fact a 3-clique). The clique number of Γ , denoted $w(\Gamma)$, is the size of the largest clique in Γ . It is hard to compute $w(\Gamma)$.


Roman Greek

$w(\Gamma)$.

Theorem: For every graph Γ , $\chi(\Gamma) \geq w(\Gamma)$.


Warning: this is not equality! For the Petersen graph P , $w(P) = 2$.

Proof: The vertices in a clique of size $w(\Gamma)$ require $w(\Gamma)$ different colors.



Dual to the clique number $\omega(\Gamma)$ we have the coclique number $\alpha(\Gamma)$ which is the maximum number of vertices in Γ , no two of which are joined. (This is $\alpha(\Gamma) = \omega(\bar{\Gamma})$ where $\bar{\Gamma}$ is the complementary graph). Cocliques are also called independent sets of vertices.

$$\text{Eg. } \alpha(P) = 4. \quad |\chi(P)| \geq \frac{10}{4} = 2.5 \Rightarrow \chi(P) \geq 3.$$



$\{1, 2, 3\}$ is a coclique which is not contained in any larger coclique; it is a maximal coclique.

A maximum coclique (i.e. a coclique of maximum size) is $\{1, 3, 7, 8\}$.

This is maximum size because P has vertex set $\{0, 3, 9, 6, 1\} \cup \{2, 7, 5, 4, 8\}$

as a union of two 5-cycles (circuits of length 5). Any set of size $\underline{4, 8}$ at least 5 vertices has either 3 on the inner 5-cycle $\{2, 7, 5, 4, 8\}$ or 3 vertices on the outer 5-cycle $\{0, 3, 9, 6, 1\}$. In either case there is an edge in that 5-cycle joining two vertices we have chosen.

Theorem $\chi(\Gamma) \geq \frac{|V|}{\alpha(\Gamma)}$ where $|V|$ = the number of vertices = the order of Γ .

Proof Let $k = \chi(\Gamma)$. Properly color the vertices $1, 2, \dots, k$ and let V_i be the subset of vertices colored i , for $i = 1, 2, \dots, k$. This gives a partition $V = V_1 \sqcup V_2 \sqcup \dots \sqcup V_k$


($V_i \cup V_j = \text{union of } V_i \text{ and } V_j$; $V_i \sqcup V_j = \text{disjoint union of } V_i \text{ and } V_j$). Each V_i is a coclique so $|V_i| \leq \alpha(\Gamma)$ so $|V| = |V_1| + |V_2| + \dots + |V_k| \leq \underbrace{\alpha(\Gamma) + \alpha(\Gamma) + \dots + \alpha(\Gamma)}_k = k\alpha(\Gamma)$ so $\chi(\Gamma) = k \geq \frac{|V|}{\alpha(\Gamma)}$. \square

March

M W F
6 8 10

Test 1 : Wed Mar 8

13 15 17 Spring Break



Using natty Software



$$G = \text{Aut } \Gamma$$






$$G = \langle (1,3)(4,5)(6,7)(8,9), (0,2)(1,4)(3,5)(6,9)(7,8) \rangle$$

$$|G| = 4$$


G has 3 orbits on the vertices: $\{0,2\}$, $\{1,3,4,5\}$, $\{6,7,8,9\}$

You can use `nauty` to test isomorphism between two graphs.





$$|\text{Aut } \overline{K'_5}| = 120$$




A graph having a Hamilton circuit is called Hamiltonian.

Looking for Hamilton paths or circuits is known to be difficult in general.

Testing whether a given graph Γ is Hamiltonian is NP-complete.

The Petersen graph P has a Hamilton path $(0, 1, 2, 3, 6, 8, 5, 7, 9)$ (a path touching each vertex exactly once) but no Hamilton circuit (ending at the same vertex where it started).

The Hamming cube $H_3 =$




does have a Hamilton circuit.

$\begin{matrix} 000 \\ 100 \\ 110 \\ 010 \\ 001 \\ 101 \\ 111 \\ 011 \\ 110 \\ 100 \end{matrix}$

"Gray code"

Every Hamming graph H_n ($n \geq 2$) has a Hamilton circuit.





Theorem: The Petersen graph P is not Hamiltonian, i.e. it does not have a Hamilton circuit / cycle.

Proof Suppose P has a Hamilton circuit. Without loss of generality this circuit contains the path $(4, 0, 1, 2)$. (This is because P has 120 automorphisms mapping any such path of length 3 to any other.) The Hamilton circuit uses two of the edges from vertex 3, so it uses either $\{3, 4\}$ or $\{2, 3\}$; so without loss of generality, it uses the edge $\{2, 3\}$.

We cannot use the edge $\{3, 4\}$ as this would complete the circuit without passing through all vertices; so we must use the edges $\{3, 6\}$ and $\{4, 5\}$. To continue the circuit from vertex 6, we have two choices: proceed through vertex 8 or vertex 9. Neither of these choices leads to a Hamilton circuit. This is a contradiction. \square

Euler paths and circuits

The Seven bridges of Königsberg



An Euler trail is a trail (repeating vertices but not edges) which uses each edge exactly once. An Euler circuit is an Euler trail that returns to its starting point.



This graph has an Euler trail. In order to have an Euler trail, a graph must have either 0 or 2 vertices of odd degree. When there are no vertices of odd degree, we have an Euler circuit.

Theorem (Euler) A graph has an Euler trail iff it is connected and it has either 0 or 2 vertices of odd degree. In the case every vertex has even degree, we have an Euler circuit/cycle.

We sometimes speak of labelled graphs and unlabelled graphs.
Eg. on the vertex set $\{1, 2, 3, 4\}$, there are $2^6 = 64$ labelled graphs
 $\binom{4}{2} = 6$ pairs of vertices.

1. .4

2. .3



There are $\binom{n}{2}$ labelled graphs on n vertices.
But many of them are isomorphic.

These are different labelled graphs but they are isomorphic.




As unlabelled graphs they are isomorphic, hence the same graph.



There are 11 unlabelled graphs of order 4 i.e. 11 isomorphism types of graphs of order 4, i.e. 11 graphs of order 4 up to isomorphism.

The Petersen graph has girth 5 (the shortest cycle has length 5). It has 15 edges. For a graph on 10 vertices, 15 edges is the maximum possible for girth 5. For a graph on 10 vertices without triangles (i.e. girth ≥ 4), what is the maximum possible number of edges?

Recall: $K_{m,n}$ =



$K_{2,8}$

16 edges



$K_{5,5}$

25 edges
girth 4
(no triangles)

$K_{9,5}$

On 9 vertices, any graph without triangles has at most 20 edges.



$K_{m,n}$ has $m \cdot n$ edges. $K_{m,n}$ is bipartite so it has no cycles of odd length. In particular it has no triangles.


Theorem (Mantel 1907)

If Γ is a graph of order n with no triangles (i.e. its girth is at least 4) then Γ has at most $\frac{n^2}{4}$ edges.

If n is even then $K_{\frac{n}{2}, \frac{n}{2}}$ attains the upper bound of $\frac{n^2}{4}$ edges. What if n is odd?

$$\left\lfloor \frac{n^2}{4} \right\rfloor = \begin{cases} \frac{n^2}{4} & \text{if } n \text{ is even, for } K_{\frac{n}{2}, \frac{n}{2}} \\ \frac{n^2-1}{4} & \text{if } n \text{ is odd, for } K_{\frac{n+1}{2}, \frac{n-1}{2}} \end{cases}$$

Proof Let Γ be a graph of order n with no triangles, $\Gamma = (V, E)$. (V is the set of vertices, E is the set of edges. For every edge $\{x, y\} \in E$, $d(x) + d(y) \leq n$.



$$d(x)-1 + d(y)-1 + d(x) + d(y) - 1 \leq n$$

Add the inequality $d(x) + d(y) \leq n$ over all edges $\{x, y\} \in E$ to get $\sum_{\{x, y\} \in E} (d(x) + d(y)) \leq ne$.

Next, count the number of triples of vertices (x, y, z) with $x \neq y \neq z$.



There are n choices for $y \in V$ and $d(y)$ choices for x , $d(y)$ choices for z , so $d(y)^2$ choices for x and z (given y). The total number of walks of length 2 is $\sum_{y \in V} d(y)^2$.

On the other hand, there are $e = |E|$ edges in Γ . For the edge $\{x, y\} \in E$, how many walks of length 2 contain this edge?

The total number of walks of length 2 is

$$\sum_{\{x, y\} \in E} (d(x) + d(y)).$$



$d(x)$ choices for z
(given the edge $\{x, y\}$)

or



$d(y)$ choices for z
(given the edge $\{x, y\}$)

a step from x to y

Therefore

$$\sum_{x \in V} d(x)^2 = \sum_{\{x,y\} \in E} (d(x) + d(y)) \leq en$$

$\underbrace{\phantom{\sum_{x \in V} d(x)^2 = \sum_{\{x,y\} \in E}}}_{\leq n}$

If $\sum_{x \in V} d(x) = 2e$, what does this tell us about $\sum_{x \in V} d(x)^2$?

Use the Cauchy-Schwarz inequality.

$$\underbrace{(d(1) + \dots + d(n))^2}_{2e} \leq n(d(1)^2 + \dots + d(n)^2)$$

$$4e^2 \leq n \sum_{x \in V} d(x)^2 = n \sum_{\{x,y\} \in E} (d(x) + d(y)) \leq n \cdot ne = n^2$$

$$\text{So } e \leq \frac{n^2}{4}.$$

Cauchy-Schwarz Inequality

Given two vectors

$$a = (a_1, a_2, \dots, a_n) \in \mathbb{R}^n$$


$$b = (b_1, b_2, \dots, b_n) \in \mathbb{R}^n,$$

$$|a \cdot b| \leq \|a\| \cdot \|b\|$$

$$\text{where } a \cdot b = a_1 b_1 + a_2 b_2 + \dots + a_n b_n$$

$$\|a\| = \sqrt{a_1^2 + a_2^2 + \dots + a_n^2}$$

$$a \cdot b = \|a\| \|b\| \cos \theta$$



$$|\cos \theta| \leq 1.$$

Special case: $b = (1, 1, 1, \dots, 1)$

$$\|b\| = \sqrt{1^2 + 1^2 + \dots + 1^2} = \sqrt{n}$$

$$a \cdot b = a_1 + a_2 + \dots + a_n$$



$$|a \cdot b| \leq \|a\| \cdot \sqrt{n}$$

$$(a_1 + \dots + a_n)^2 \leq n(a_1^2 + \dots + a_n^2)$$

Proof of Cauchy-Schwarz: Fix $a, b \in \mathbb{R}^n$. Consider

$$f(t) = \|a - tb\|^2 = (a - tb) \cdot (a - tb) = a \cdot a - tb \cdot a - ta \cdot b + t^2 b \cdot b = \|a\|^2 - 2t(a \cdot b) + t^2 \|b\|^2.$$

of course $f(t) \geq 0$ for all t .



The discriminant

$$(-2a \cdot b)^2 - 4\|a\|^2\|b\|^2 \leq 0$$

$$4(a \cdot b)^2 \leq 4\|a\|^2\|b\|^2$$

$$(a \cdot b)^2 \leq \|a\|^2\|b\|^2.$$


The disc. of at^2+bt+c
is b^2-4ac

Show me a graph Γ of order 5 such that neither Γ nor $\bar{\Gamma}$ has a triangle.
 (i.e. both Γ and $\bar{\Gamma}$ have girth ≥ 4). Recall: $\bar{\Gamma}$ is the complement of Γ .

$$\Gamma = \text{pentagon} \quad \bar{\Gamma} = \text{star} \cong \Gamma \quad (\text{a } 5\text{-cycle}). \text{ Both } \Gamma \text{ and } \bar{\Gamma} \text{ have girth 5.}$$


Show me a graph Γ of order 6 such that neither Γ nor $\bar{\Gamma}$ have a triangle.
 There is no such graph. Why not?

Color the edges of K_5 with 2 colors red, blue. To avoid a monochromatic triangle (all red or all blue):



Theorem If we color the edges of K_6 red and blue,
 then there is either a red triangle or a blue triangle.

Proof Consider a vertex v .



Now x, y, z form a triangle.

If any edge of this triangle is blue
 then together with the edges to v we have
 a blue triangle. Otherwise all edges
 of the triangle $\{x, y, z\}$ are red. \square

There are five
 edges from v
 So by the
 Pigeonhole
 Principle, at
 least three of
 them are the
 same color, say
 $\{v, x\}, \{v, y\}, \{v, z\}$
 are blue.

Ramsey Theorem: Let r, s be positive integers. There is a number $R(r, s)$ such that for all $n \geq R(r, s)$, every 2-coloring of the edges of K_n has either a red r -clique or a blue s -clique. For $n < R(r, s)$, there exists a coloring of the edges of K_n with 2 colors (red, blue) having no red r -clique and no blue s -clique.

Values / known bounding ranges for Ramsey numbers $R(r, s)$ (sequence A212954 in the OEIS)

$$R(3, 3) = 6.$$

$$R(2, 3) = 3$$




$$43 \leq R(5, 5) \leq 48$$


$R(r, s)$ values
are Ramsey
numbers.

(Extremal graph
theory)


$r \backslash s$	1	2	3	4	5	6	7	8	9	10
1	1	1	1	1	1	1	1	1	1	1
2		2	3	4	5	6	7	8	9	10
3			6	9	14	18	23	28	36	40–42
4				18	25 ^[9]	36–40	49–58	59 ^[13] –79	73–106	92–136
5					43–48	58–85	80–133	101–194	133–282	149 ^[13] –381
6						102–161	115 ^[13] –273	134 ^[13] –427	183–656	204–949
7							205–497	219–840	252–1379	292–2134
8								282–1532	329–2683	343–4432
9									565–6588	581–12677
10									798–23556	



is a subgraph of Γ



is an induced subgraph.



is not a
subgraph.