



Fields

Book III

We have been talking about number fields: finite extensions $E \supseteq \mathbb{Q}$ i.e. $[E:\mathbb{Q}] = n < \infty$.
(Some are Galois i.e. $G = \text{Aut } E$ satisfies $|G| = n$; but in general $|G| \leq n$.)

Back to basics:

In a field F , if $\underbrace{1+1+\dots+1}_{n \geq 1} = 0$ then the smallest n for which this occurs is the characteristic of F .

If F has characteristic $n > 0$ then n must be prime. If $n = ab$, $a, b \geq 1$ then

$$\underbrace{(1+1+\dots+1)}_a \underbrace{(1+1+\dots+1)}_b = \underbrace{1+1+\dots+1}_{n=ab} = 0$$

By minimality of n , n is prime.

If $\underbrace{1+1+\dots+1}_n \neq 0$ for any $n \geq 1$, then we say n has characteristic 0.

Given a field F , $\text{char } F =$ characteristic of F is either 0 or p (some prime p).

• If $\text{char } F = p$ then $F \supseteq \mathbb{F}_p =$ field of order p ($\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z} = \{0, 1, 2, \dots, p-1\} =$ "integers mod p ").

eg. $\mathbb{F}_p, \mathbb{F}_{p^2}, \mathbb{F}_{p^3}, \mathbb{F}_{p^4}, \dots, \mathbb{F}_p(x) = \{ \text{all rational functions in } x \text{ with coefficients in } \mathbb{F}_p \}, \dots$

• If $\text{char } F = 0$ then $F \supseteq \mathbb{Q}$. Eg. $\mathbb{R}, \mathbb{C}, \mathbb{Q}$, number fields, $A = \{ \text{algebraic numbers} \} \subset \mathbb{C}$
eg. $\mathbb{Q}[\sqrt{2}]$

In either case F has a unique smallest subfield, either \mathbb{F}_p or \mathbb{Q} , called the prime subfield of F .

All fields of characteristic 0 are infinite. (They are extensions of \mathbb{Q} , hence vector spaces over \mathbb{Q} .)

If $E \supseteq F$ is a field extension (i.e. E, F are fields with F a subfield of E) then E is a vector space over F . The dimension of this vector space is the degree $[E:F]$ of this extension eg.

$$[\mathbb{C}:\mathbb{R}] = 2$$

$\{1, i\}$ basis

$$[\mathbb{R}:\mathbb{Q}] = \infty$$

$1, \sqrt{2}, \sqrt{3}, \sqrt{5}, \sqrt{6}, \sqrt{7}, \sqrt{10}, \sqrt{11}, \dots$
are lin. indep.

$$[\mathbb{C}:\mathbb{Q}] = \underbrace{[\mathbb{C}:\mathbb{R}]}_2 \underbrace{[\mathbb{R}:\mathbb{Q}]}_{\infty} = \infty$$

For fields of characteristic a prime p , some are finite, some are infinite.

Given p prime and $k \geq 1$ (positive integer), there is a unique field of order $q = p^k$ (up to isomorphism)

Finite fields: $\mathbb{F}_2, \mathbb{F}_3, \mathbb{F}_4, \mathbb{F}_5, \mathbb{F}_7, \mathbb{F}_8, \mathbb{F}_9, \mathbb{F}_{11}, \mathbb{F}_{13}, \mathbb{F}_{16}, \mathbb{F}_{17}, \dots$

$$\mathbb{F}_4 = \{0, 1, \alpha, \beta\}$$

+	0	1	α	β
0	0	1	α	β
1	1	0	β	α
α	α	β	0	1
β	β	α	1	0

$$\alpha + \alpha = (1+1)\alpha = 0\alpha = 0$$

×	0	1	α	β
0	0	0	0	0
1	0	1	α	β
α	0	α	β	1
β	0	β	1	α

$$\text{char } \mathbb{F}_4 = 2.$$

$\mathbb{F}_4 \supset \mathbb{F}_2$ of degree $[\mathbb{F}_4:\mathbb{F}_2] = 2$

with basis $1, \alpha$

$$\mathbb{F}_4 = \{a \cdot 1 + b\alpha : a, b \in \mathbb{F}_2\}$$

$$= \{0, 1, \alpha, 1+\alpha\} \quad \text{where } \alpha^2 = \alpha + 1.$$

$$= \{0, 1, \alpha, \alpha^2\} \quad \beta$$

$$\mathbb{F}_4 = \mathbb{F}_2[\alpha]$$

The minimal poly. of α over \mathbb{F}_2 is $x^2 + x + 1$.

Irreducible polynomials over $\mathbb{F}_2 = \{0, 1\}$

degree 1: $x, x+1$ (both irreducible)

degree 2: $x^2, x^2+1, x^2+x, x^2+x+1$
 $\underbrace{x \cdot x \quad (x+1)(x+1) \quad x(x+1)}_{\text{reducible}}$ irreducible

degree 3: $x^3 = x \cdot x \cdot x$
 $x^3+1 = (x+1)(x^2+x+1)$
 $x^3+x = x \cdot (x+1)^2$
 x^3+x+1 irreducible
 $x^3+x^2 = x \cdot x \cdot (x+1)$
 x^3+x^2+1 irreducible
 $x^3+x^2+x = x(x^2+x+1)$
 $x^3+x^2+x+1 = (x+1)^3$

In general the nonzero elements of \mathbb{F}_q form a cyclic group of order $q-1$.

There is only one finite field of each order $q=p^k$ (p prime, $k \geq 1$) up to isomorphism.

If \mathbb{F}_q is a finite field then it must have $\text{char } \mathbb{F}_q = p$ for some prime p

$$|\mathbb{F}_q| = q < \infty$$

So \mathbb{F}_q is an extension $\mathbb{F}_q \supseteq \mathbb{F}_p$ hence a vector space of some dimension k .
 Let $\alpha_1, \dots, \alpha_k$ be a basis for \mathbb{F}_q over \mathbb{F}_p ie. $\mathbb{F}_q = \{a_1\alpha_1 + a_2\alpha_2 + \dots + a_k\alpha_k : a_1, \dots, a_k \in \mathbb{F}_p\}$.

$$q = |\mathbb{F}_q| = p^k$$

There are 2^n polynomials of degree $n : x^n + c_{n-1}x^{n-1} + \dots + c_1x + c_0$ and they are all monic. $c_0, c_1, \dots, c_{n-1} \in \mathbb{F}_2$

Let α be a root of x^2+x+1 . The other root is $\alpha+1$.

$$\alpha^2 + \alpha + 1 = 0 \Rightarrow \alpha^2 = -\alpha - 1 = \alpha + 1$$

Note: The roots of $ax^2+bx+c=0$ are $\frac{-b \pm \sqrt{b^2-4ac}}{2a}$ except in characteristic 2.

$\mathbb{F}_8 = \mathbb{F}_2[\gamma]$ where γ is a root of x^3+x+1
 $= \{a + b\gamma + c\gamma^2 : a, b, c \in \mathbb{F}_2\}$
 $= \{0, 1, \gamma, \gamma+1, \gamma^2, \gamma^2+1, \gamma^2+\gamma, \gamma^2+\gamma+1\}$

 $\underbrace{\quad}_{\gamma^3} \quad \underbrace{\quad}_{\gamma^6} \quad \underbrace{\quad}_{\gamma^4} \quad \underbrace{\quad}_{\gamma^5}$

$$\text{ie. } \gamma^3 = \gamma + 1$$

$$\gamma^0 = 1$$

$$\gamma^1 = \gamma$$

$$\gamma^2 = \gamma^2$$

$$\gamma^3 = \gamma + 1$$

$$\gamma^4 = \gamma^2 + \gamma$$

$$\gamma^5 = \gamma^3 + \gamma^2 = \gamma^2 + \gamma + 1$$

$$\gamma^6 = \gamma^3 + \gamma^2 + \gamma = (\gamma + 1) + \gamma^2 + \gamma$$

$$= \gamma^2 + 1$$

$$\gamma^7 = \gamma^3 + \gamma = (\gamma + 1) + \gamma = 1$$

x^3+x+1 has three roots in \mathbb{F}_8 :

$$\gamma, \gamma^2, \gamma^4$$

x^3+x^2+1 has three roots in \mathbb{F}_8 :

$$\gamma^3, \gamma^5, \gamma^6 = \gamma^7$$

$$\mathbb{F}_9 = \mathbb{F}_3[i] \quad \text{compare: } \mathbb{C} = \mathbb{R}[i],$$

$$= \{a+bi : a, b \in \mathbb{F}_3\}$$

$$= \{0, 1, 2, i, 1+i, 2+i, 2i, 1+2i, 2+2i\}$$

$$\begin{array}{ccccccc} \theta^0 & \theta^1 & \theta^2 & \theta^3 & \theta^4 & \theta^5 & \theta^6 \\ \theta^3 & \theta^4 & \theta^5 & \theta^6 & \theta^7 & \theta^8 & \theta^9 \end{array}$$

$$\mathbb{Q}[i] \supset \mathbb{Q}, \quad i = \sqrt{-1}. \quad \{1, i\} \text{ is a basis of the extension}$$

$$\mathbb{Q}[\sqrt{2}] \supset \mathbb{Q} \quad \text{in each case.}$$

$$i = \sqrt{-1} = \sqrt{2} \quad \mathbb{F}_9 = \mathbb{F}_3[i] = \mathbb{F}_3[\sqrt{2}]$$

$$\theta^0 = 1$$

$$\theta^1 = \theta = 1+i$$

$$\theta^2 = (1+i)^2 = 1+2i+i^2 = 2i$$

$$\theta^3 = \frac{2i(1+i)}{\theta} = -2+2i = 1+2i$$

$$\theta^4 = \theta^3 \cdot \theta = (1+2i)(1+i) = 1-2 = -1 = 2$$

$$\theta^5 = \theta^4 \cdot \theta = -\theta = 2\theta = 2+2i$$

$$\theta^6 = \theta^5 \cdot \theta = -\theta^2$$

$$\theta^7 = \theta^6 \cdot \theta = -\theta^3$$

$$\theta^8 = \theta^7 \cdot \theta = -\theta^4$$