



# Fields

Book I

# Fields

Let  $F$  be a set containing distinct elements called 0 and 1 (thus  $0 \neq 1$ ). Suppose addition, subtraction, multiplication and division are defined for all elements of  $F$  (except division by 0 is not defined).

Thus  $a + b, a - b, ab, \frac{a}{d} \in F$  whenever  $a, b, d \in F$  and  $d \neq 0$ .

Define  $-a = 0 - a$ .

If the following properties are satisfied by *all* elements  $a, b, c, d \in F$  with  $d \neq 0$ , then  $F$  is a **field**.

$$a + b = b + a$$

$$a + 0 = a$$

$$a + (-a) = 0$$

$$a + (-b) = a - b$$

$$a + (b + c) = (a + b) + c$$

$$a(bc) = (ab)c$$

$$a(b + c) = ab + ac$$

$$ab = ba$$

$$1a = a$$

$$\frac{a}{d}d = a$$

$\mathbb{Q}^{2 \times 2} = \{2 \times 2 \text{ matrices over } \mathbb{Q}\} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{Q} \right\}$  is not a field.

$$0 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \quad I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \text{ identity}$$

$$A + 0 = A, \quad AI = A = IA$$

$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$  has no inverse.  $A \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = I$  has no solution for  $A$ .

Moreover,  $AB \neq BA$  in general.

$\mathbb{Q}^{2 \times 2}$  is a (non-commutative) ring with identity.

It has a subring  $D = \left\{ \begin{bmatrix} a & 0 \\ 0 & d \end{bmatrix} : a, d \in \mathbb{Q} \right\}$  is a commutative subring with identity.

But  $D$  is not a field since it has non-invertible elements.

$D$  has zero divisors:  $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ . A field can never have zero divisors.

(If  $d$  is a zero divisor then  $cd = 0$  where  $c, d \neq 0$  so  $\left(\frac{c}{d}\right)d = c \neq 0$ , contradiction)

For a commutative ring  $R$  with identity, being able to divide is stronger than having no zero divisors.

An example of a commutative ring with identity having no zero divisors but not a field (division fails in general) is  $\mathbb{Z}$

Eg.  $F = \left\{ \begin{bmatrix} a & b \\ 2b & a \end{bmatrix} : a, b \in \mathbb{Q} \right\} \subset \mathbb{Q}^{2 \times 2}$  is a subring, containing  $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ .

If  $\begin{bmatrix} a & b \\ 2b & a \end{bmatrix} \neq \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$  then  $\begin{bmatrix} a & b \\ 2b & a \end{bmatrix}^{-1} = \frac{1}{a^2 - 2b^2} \begin{bmatrix} a & -b \\ -2b & a \end{bmatrix}$  (Note:  $a^2 - 2b^2 \neq 0$  since  $\sqrt{2} \notin \mathbb{Q}$ )

Why is  $F$  a commutative subring? Elements of  $F$  have the form

$\begin{bmatrix} a & b \\ 2b & a \end{bmatrix} = aI + bS$  where  $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ ,  $S = \begin{bmatrix} 0 & 1 \\ 2 & 0 \end{bmatrix}$  so  $F = \{aI + bS : a, b \in \mathbb{Q}\}$  is the span of  $\{I, S\}$  in  $\mathbb{Q}^{2 \times 2}$  ( $F$  is a 2-dimensional subspace of  $\mathbb{Q}^{2 \times 2}$ , a 4-dimensional vector space).

$$\begin{aligned} \begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} &= \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \\ &= \begin{bmatrix} \frac{d}{ad-bc} & \frac{-b}{ad-bc} \\ \frac{-c}{ad-bc} & \frac{a}{ad-bc} \end{bmatrix} \end{aligned}$$

$$(aI + bS)(cI + dS) = acI + (ad + bc)S + bdS^2 = (cI + dS)(aI + bS), \quad S^2 = \begin{bmatrix} 0 & 1 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 2 & 0 \end{bmatrix} = 2I$$

$$= (ac + 2bd)I + (ad + bc)S$$


---

Compare:  $K = \mathbb{Q}[\sqrt{2}] = \{a + b\sqrt{2} : a, b \in \mathbb{Q}\}$  is a field.

$$(a + b\sqrt{2}) + (c + d\sqrt{2}) = (a + c) + (b + d)\sqrt{2}$$

$$(a + b\sqrt{2})(c + d\sqrt{2}) = ac + (ad + bc)\sqrt{2} + 2bd = (ac + 2bd) + (ad + bc)\sqrt{2}$$

Note:  $F \cong K$  (they are isomorphic)

An explicit isomorphism  $\phi: K \rightarrow F$  is given by  $\phi(a + b\sqrt{2}) = \begin{bmatrix} a & b \\ 2b & a \end{bmatrix} = aI + bS$ .

$\phi$  is bijective

$$\phi(x + y) = \phi(x) + \phi(y)$$

$$\phi(xy) = \phi(x)\phi(y)$$


---

Similarly  $\left\{ \begin{bmatrix} a & b \\ -b & a \end{bmatrix} : a, b \in \mathbb{R} \right\} \subset \mathbb{R}^{2 \times 2}$  is a subring isomorphic to  $\mathbb{C}$ .

An isomorphism  $\mathbb{C} \rightarrow \left\{ \begin{bmatrix} a & b \\ -b & a \end{bmatrix} : a, b \in \mathbb{R} \right\}$  is  $a + bi \mapsto \begin{bmatrix} a & b \\ -b & a \end{bmatrix}$  ( $a, b \in \mathbb{R}$ ).

$$\mathbb{Q}[\sqrt{2}] = \{ a + b\sqrt{2} : a, b \in \mathbb{Q} \}$$

$$\alpha = 5 + 3\sqrt{2}, \quad \beta = 7 - \sqrt{2}$$

$$\alpha + \beta = 12 + 2\sqrt{2}$$

$$\alpha - \beta = -2 + 4\sqrt{2}$$

$$\alpha\beta = (5 + 3\sqrt{2})(7 - \sqrt{2}) = 35 - 5\sqrt{2} + 21\sqrt{2} - 6 = 29 + 16\sqrt{2}$$

$$\frac{\alpha}{\beta} = \frac{5 + 3\sqrt{2}}{7 - \sqrt{2}} = \frac{5 + 3\sqrt{2}}{7 - \sqrt{2}} \cdot \frac{7 + \sqrt{2}}{7 + \sqrt{2}} = \frac{35 + 5\sqrt{2} + 21\sqrt{2} + 6}{47} = \frac{41 + 26\sqrt{2}}{47} = \frac{41}{47} + \frac{26}{47}\sqrt{2}$$

Alternatively,  $\frac{\alpha}{\beta} = \alpha\beta^{-1}$

in matrix representation:  $\begin{bmatrix} 5 & 3 \\ 6 & 5 \end{bmatrix} \cdot \frac{1}{47} \begin{bmatrix} 7 & 1 \\ 2 & 7 \end{bmatrix} = \frac{1}{47} \begin{bmatrix} 41 & 26 \\ 52 & 41 \end{bmatrix}$

$$\beta \mapsto \begin{bmatrix} 7 & -1 \\ -2 & 7 \end{bmatrix}$$

$$\beta^{-1} \mapsto \frac{1}{47} \begin{bmatrix} 7 & 1 \\ 2 & 7 \end{bmatrix}$$

Similar:  $\mathbb{Q}[\sqrt[3]{2}] = \mathbb{Q}[\theta]$ ,  $\theta = \sqrt[3]{2}$ .

$\{ a + b\theta : a, b \in \mathbb{Q} \}$  is not a field, not even a ring, since it's not closed under multiplication.

$\mathbb{Q}[\theta] = \{ a + b\theta + c\theta^2 : a, b, c \in \mathbb{Q} \}$  is a field.

$$\alpha = 5 + 3\theta$$

$$\beta = 7 - \theta$$

$$\alpha + \beta = 12 + 2\theta$$

$$\alpha - \beta = -2 + 4\theta$$

$$\alpha\beta = (5 + 3\theta)(7 - \theta) = 35 - 5\theta + 21\theta - 3\theta^2 = 35 + 16\theta - 3\theta^2$$

$$\theta^3 = 2$$

$$\theta^4 = 2\theta$$

$$\theta^5 = 2\theta^2$$

$$\theta^6 = 4$$

$$\frac{\alpha}{\beta} = \frac{5+3\theta}{7-\theta} = \frac{a}{1} + \frac{b}{1}\theta + \frac{c}{1}\theta^2 = \frac{251}{341} + \frac{182}{341}\theta + \frac{26}{341}\theta^2 = \frac{1}{341}(251 + 182\theta + 26\theta^2)$$

$$\theta^3 = 2$$

$$\theta^3 - 2 = 0$$

$$\theta = \sqrt[3]{2}$$


rational coefficients  
 $a, b, c \in \mathbb{Q}$

$\theta$  is a root of  $x^3 - 2 = (x - \theta)(x^2 + \theta x + \theta^2)$

$$5 + 3\theta = (a + b\theta + c\theta^2)(7 - \theta)$$

$$= 7a + (7b - a)\theta + (7c - b)\theta^2 - 2c$$

$$= (7a - 2c) + (7b - a)\theta + (7c - b)\theta^2$$



Hopefully

$$\begin{aligned} 7a - 2c &= 5 \\ -a + 7b &= 3 \\ -b + 7c &= 0 \end{aligned}$$

$$\left[ \begin{array}{ccc|c} 7 & 0 & -2 & 5 \\ -1 & 7 & 0 & 3 \\ 0 & -1 & 7 & 0 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 0 & 19 & -2 & 26 \\ -1 & 7 & 0 & 3 \\ 0 & -1 & 7 & 0 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & -7 & 0 & -3 \\ 0 & 19 & -2 & 26 \\ 0 & 1 & -7 & 0 \end{array} \right]$$

$$\sim \left[ \begin{array}{ccc|c} 1 & -7 & 0 & -3 \\ 0 & 1 & -7 & 0 \\ 0 & 19 & -2 & 26 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 0 & -49 & -3 \\ 0 & 1 & -7 & 0 \\ 0 & 0 & 341 & 26 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 0 & -49 & -3 \\ 0 & 1 & -7 & 0 \\ 0 & 0 & 1 & \frac{26}{341} \end{array} \right]$$

$$\sim \left[ \begin{array}{ccc|c} 1 & 0 & 0 & \frac{251}{341} \\ 0 & 1 & 0 & \frac{182}{341} \\ 0 & 0 & 1 & \frac{26}{341} \end{array} \right]$$

$$\begin{array}{r} 26 \\ 7 \\ \hline 182 \end{array}$$

$$\begin{array}{r} 49 \\ 26 \\ \hline 244 \\ 98 \\ \hline 1274 \end{array}$$

$$\begin{array}{r} 341 \\ 3 \\ \hline 1023 \end{array}$$

$$= -3 + 49 \cdot \frac{26}{341}$$

$$= -3 + \frac{1274}{341}$$

$$= \frac{-1023 + 1274}{341} = \frac{251}{341}$$

$$\text{Check: } \frac{1}{341}(251 + 182\theta + 26\theta^2)(7 - \theta) = \frac{1}{341}(1757 + 1023\theta + 0\theta^2 - 52)$$

$$= \frac{1}{341}(1705 + 1023\theta)$$

$$= 5 + 3\theta \quad \checkmark$$

$\mathbb{Q}[\theta]$  is a cubic field extension of  $\mathbb{Q}$ : it is a 3-dimensional vector space over  $\mathbb{Q}$ , with basis  $1, \theta, \theta^2$ .

Alternatively, use  $3 \times 3$  matrices to represent elements of  $\mathbb{Q}[\theta]$ .

Take  $T = \begin{bmatrix} 0 & 0 & 2 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$  to represent  $\theta$ .  $T^3 = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 0 & 2 \\ 0 & 0 & 2 \end{bmatrix} = 2I$

$$E = \left\{ aI + bT + cT^2 : a, b, c \in \mathbb{Q} \right\} = \left\{ \begin{bmatrix} a & 2c & 2b \\ b & a & 2c \\ c & b & a \end{bmatrix} : a, b, c \in \mathbb{Q} \right\} \subset \mathbb{Q}^{3 \times 3}$$

noncommutative  
ring with identity  
having zero divisors

$\mathbb{Q}[\theta] \cong E$  via the isomorphism

$$a + b\theta + c\theta^2 \mapsto aI + bT + cT^2$$

This subring is  
a field.

Are there any fields "between"  $\mathbb{Q}$  and  $\mathbb{Q}[\sqrt{2}]$ , or between  $\mathbb{Q}$  and  $\mathbb{Q}[\theta]$ ?

Are there any fields "between"  $\mathbb{R}$  and  $\mathbb{C}$ ?

Suppose  $\mathbb{R} \subset F \subset \mathbb{C}$  is a tower of fields ( $F$  is a subfield of  $\mathbb{C}$  and  $\mathbb{R}$  is a subfield of  $F$ ).  $\subseteq$  vs  $\subset$  'C' always means strict containment in this course.

Since  $F \supset \mathbb{R}$ , there exists  $\alpha \in F$ ,  $\alpha \notin \mathbb{R}$ . Then  $\alpha, 1$  are linearly independent over  $\mathbb{R}$ , i.e.  $\alpha \neq a \cdot 1$  for any  $a \in \mathbb{R}$ . However  $\mathbb{C}$  is 2-dimensional over  $\mathbb{R}$  with basis  $1, i$  (every complex number is uniquely expressible as  $z = a \cdot 1 + b \cdot i$  with  $a, b \in \mathbb{R}$ ). So  $1, \alpha$  is a basis for  $F$ . So  $F = \mathbb{C}$ .

Is there any field extension  $\mathbb{C} \subset F$  with  $F$  2-dimensional over  $\mathbb{C}$ ?  
 No, but there do exist fields  $F \supset \mathbb{C}$  which are infinite-dimensional extensions.

Consider the ring  $\mathbb{C}[x] = \{ \text{polynomials in } x \text{ with complex coefficients} \}$

This is a ring but not quite a field eg.  
 $= \{ a_0 + a_1x + a_2x^2 + \dots + a_nx^n : a_i \in \mathbb{C}, n \geq 0 \}$

$$\frac{5+7x+ix^2}{3-(1+i)x+43x^2} \notin \mathbb{C}[x]$$

$\mathbb{C}(x) =$  field of fractions of  $\mathbb{C}[x]$

$=$  field of rational functions in  $x$  with complex coefficients

Just like constructing  $\mathbb{Q}$  from  $\mathbb{Z}$ .

Another example of this: We'll construct a countably infinite subfield of  $\mathbb{R}$  containing  $\pi$ .

This contains the subring  $\mathbb{Q}[\pi] = \{ a_0 + a_1\pi + a_2\pi^2 + \dots + a_n\pi^n : n \geq 0, a_i \in \mathbb{Q} \}$

$\pi \in \mathbb{Q}[\pi]$  has no (multiplicative) inverse in  $\mathbb{Q}[\pi]$  since if

$$1 = \pi (a_0 + a_1\pi + a_2\pi^2 + \dots + a_n\pi^n) \quad a_i \in \mathbb{Q}, n \geq 0,$$

a contradiction since  $\pi$  is transcendental. ( $\pi$  would be a root of a nonzero polynomial  $a_nx^{n+1} + a_{n-1}x^n + \dots + a_2x^3 + a_1x^2 + a_0x - 1$ )  
 (Lindemann 1800's)

$\mathbb{Q}(\pi) = \{ \frac{a}{b} : a, b \in \mathbb{Q}[\pi], b \neq 0 \}$  is the field of quotients of the ring  $\mathbb{Q}[\pi]$

$\mathbb{Q}(\sqrt{2}) = \{ \frac{a}{b} : a, b \in \mathbb{Q}[\sqrt{2}], b \neq 0 \} = \mathbb{Q}[\sqrt{2}]$  is already a field.  $\sqrt{2}$  is algebraic: it is a root of a nonzero poly.  $x^2 - 2 \in \mathbb{Q}[x]$



Every  $\alpha \in \mathbb{C}$  is either algebraic or transcendental, never both.



$\mathbb{Q} \subset \mathbb{R} \subset \mathbb{C}$   
 countable      uncountable      uncountable

$\mathbb{Q} \subset \mathbb{A} \subset \mathbb{C}$   
 countable      uncountable.

$A = \{\text{algebraic numbers}\}$



Elements of  $\mathbb{Q}(\pi) \subset \mathbb{R}$  look like

$$\frac{53.8\pi^2 - 17\pi + \frac{5}{7}}{12\pi^2 + 119\pi + \frac{103}{648}}$$

Compare:  $\mathbb{Q}(e) \subset \mathbb{R}$ , another countable subfield of  $\mathbb{R}$ .  
 Actually  $\mathbb{Q}(e) \cong \mathbb{Q}(\pi) \cong \mathbb{Q}(x)$  (x being an indeterminate)

$\mathbb{Q}[x]$  is a countably infinite ring  
 so  $\mathbb{Q}(\pi)$  is a countably infinite field.

An isomorphism is  $f(e) \mapsto f(\pi)$  where  $f(x) \in \mathbb{Q}(x)$ .  
 ie. an abstract general symbol

$\mathbb{Q}(x) \rightarrow \mathbb{Q}(\pi)$  evaluation

$\mathbb{Q}(x) \rightarrow \mathbb{Q}(e)$

$\mathbb{Q}(x) \rightarrow \mathbb{Q}(\sqrt{2})$  doesn't quite work eg. the image of  $\frac{x^3 + 7x^2 - 3}{x^2 - 2} \in \mathbb{Q}(x)$  is undefined; you can't evaluate this at  $\sqrt{2}$ .

$\mathbb{Q}[x] \rightarrow \mathbb{Q}[\pi]$

$\mathbb{Q}[x] \rightarrow \mathbb{Q}[e]$

$\mathbb{Q}[x] \rightarrow \mathbb{Q}[\sqrt{2}]$

} all well-defined ring homomorphisms.

But the evaluation maps at  $\pi, e, \sqrt{2}, \dots$

-3 -2 -1 0 1 2

If  $\phi: R \rightarrow S$  where  $R, S$  are rings, we say  $\phi$  is a ring homomorphism if

$$\left. \begin{aligned} \phi(a+b) &= \phi(a) + \phi(b) \\ \phi(ab) &= \phi(a)\phi(b) \end{aligned} \right\} \text{ for all } a, b \in R$$

We don't necessarily require  $\phi(1) = 1$ ; and in general the rings  $R, S$  may not have identity.

If  $R, S$  are rings with identity ( $1_R \in R, 1_S \in S$ ) we might consider only homomorphisms of rings with identity i.e.  $\phi(1_R) = \phi(1_S)$ .

\* Suppose  $F, K$  are fields. If  $\phi: F \rightarrow K$  is a ring homomorphism then either (trivial)

(i)  $\phi(F) = \{0\}$  i.e.  $\phi(a) = 0$  for all  $a \in F$ , or

(ii)  $\phi$  is one-to-one i.e.  $\phi(F) \subseteq K$  is a subfield isomorphic to  $F$ .

Any homomorphism  $\mathbb{Q}[x] \rightarrow \mathbb{R}$  is either trivial or it has the form  $\mathbb{Q}[x] \rightarrow \mathbb{Q}(a), f(x) \mapsto f(a)$  is an evaluation at some transcendental number  $a \in \mathbb{R}$ .

We have ring homomorphisms  $\mathbb{Q}[x] \rightarrow \mathbb{C}^{n \times n}$  ( $n \times n$  complex matrices) where we evaluate at a matrix  $A \in \mathbb{C}^{n \times n}$ , i.e.  $f(x) \mapsto f(A)$

$$\frac{47}{3}x^2 + \frac{18}{11}x - \frac{11}{7} \mapsto \frac{47}{3}A^2 + \frac{18}{11}A - \frac{11}{7}I$$

(\*) In a field  $F$ , every ideal is either  $\{0\}$  or  $F$ .

An automorphism of a field  $F$  is an isomorphism  $\phi: F \rightarrow F$ . Eg. bijective with

(i) Automorphisms of  $\mathbb{Q}[\sqrt{2}]$ ? We want  $\phi: \mathbb{Q}[\sqrt{2}] \rightarrow \mathbb{Q}[\sqrt{2}]$

$$\phi(a+b) = \phi(a) + \phi(b), \quad \phi(ab) = \phi(a)\phi(b).$$

• The identity  $\phi(x) = x$  for all  $x \in \mathbb{Q}[\sqrt{2}]$

• Conjugation  $\phi(a+b\sqrt{2}) = a-b\sqrt{2}$  for all  $a, b \in \mathbb{Q}$ . (This is algebraic conjugation, not complex conjugation).

These are the only automorphisms of  $\mathbb{Q}[\sqrt{2}]$ .

If  $\phi: F \rightarrow F$  is any automorphism of a field  $F$  then

$$\phi(0) = \phi(0+0) = \phi(0) + \phi(0) \Rightarrow \phi(0) = 0$$

$$\phi(1) = \phi(1 \cdot 1) = \phi(1) \cdot \phi(1) \text{ where } \phi(1) \neq 0 \text{ since } \phi \text{ is one-to-one. Multiply both sides}$$

by  $\phi(1)^{-1}$  to get  $\phi(1) = 1$ .

If  $m, n \in \mathbb{Z}$  with  $n \neq 0$ ,

$$\phi\left(n \cdot \frac{m}{n}\right) = \phi(m) = m$$

$$\phi\left(\frac{m}{n}\right) \phi(n) \Rightarrow \phi\left(\frac{m}{n}\right) = \frac{m}{n}$$

So  $\phi(x) = x$  for all  $x \in \mathbb{Q}$ .

$$\phi(2) = \phi(1+1) = \phi(1) + \phi(1) = 1+1 = 2$$

$$\phi(3) = \phi(2+1) = \phi(2) + \phi(1) = 2+1 = 3$$

So  $3 + (-3) = 0$

$$\phi(3) + \phi(-3) = \phi(0) = 0$$

$$\phi(3) = 3$$

$$\phi(-3) = -3$$

$$\phi(\sqrt{2})^2 = \phi(\sqrt{2}^2) = \phi(2) = 2 \Rightarrow \phi(\sqrt{2}) = \pm\sqrt{2}$$

for all  $a, b \in \mathbb{R}$

$$\text{If } \phi(\sqrt{2}) = \sqrt{2} \text{ then } \phi(a+b\sqrt{2}) = \phi(a) + \phi(b)\phi(\sqrt{2}) = a + b\sqrt{2}$$

$$\text{If } \phi(\sqrt{2}) = -\sqrt{2} \text{ then } \phi(a+b\sqrt{2}) = \phi(a) + \phi(b)\phi(\sqrt{2}) = a + b(-\sqrt{2}) = a - b\sqrt{2}$$

If  $F$  is any field then  $\text{Aut } F = \{\text{all automorphisms of } F\}$  is a group under composition. Its identity is  $\iota$  where  $\iota: F \rightarrow F$ ,  $\iota(x) = x$  for all  $x \in F$  (the identity map).

$\text{Aut } \mathbb{Q} = \{\iota\}$  is trivial.

$\text{Aut } \mathbb{R} = \{\iota\}$  is trivial but why?

$\mathbb{Q}[\sqrt{2}] \subset \mathbb{R}$  has two automorphisms.

$\text{Aut } \mathbb{Q}[\sqrt{2}]$  is a group of order 2.


$\text{Aut } \mathbb{C}$  contains  $\iota$  and  $\tau = \text{complex conjugation}$ ,

But  $\text{Aut } \mathbb{C}$  is uncountable.  $\mathbb{C}$  has uncountably many automorphisms.


The only continuous automorphisms of  $\mathbb{C}$  are  $\iota, \tau$ .

$$\tau(a+bi) = a-bi \text{ for all } a, b \in \mathbb{R}.$$

The conjugation  $\phi \in \text{Aut } \mathbb{Q}[\sqrt{2}]$  defined by  $\phi(a+b\sqrt{2}) = a-b\sqrt{2}$  ( $a, b \in \mathbb{Q}$ ) is badly discontinuous



$l: \mathbb{Q}[\sqrt{2}] \rightarrow \mathbb{Q}[\sqrt{2}]$   
 $l$  is continuous



graph of  $\phi$   
 $\phi$  is badly discontinuous.


$\mathbb{R}(x) = \{ \text{rational functions of } x \text{ with real coefficients} \}$  is a field.  
 Can we replace "rational functions" with "functions" or "continuous functions"  $\mathbb{R} \rightarrow \mathbb{R}$  ?

$\{ \text{functions } \mathbb{R} \rightarrow \mathbb{R} \}$


$\{ \text{continuous functions } \mathbb{R} \rightarrow \mathbb{R} \}$

are rings with zero divisors so they are not fields.

Commutative rings with identity under pointwise multiplication.



$$f_1(x) = \begin{cases} x, & \text{if } x \geq 0 \\ 0, & \text{if } x < 0 \end{cases}$$



$$f_2(x) = \begin{cases} 0, & \text{if } x \geq 0 \\ x, & \text{if } x < 0 \end{cases}$$

$f_1 f_2 = 0$  but  $f_1, f_2$  are nonzero functions.

How do we check that  $f(x) \in \mathbb{Q}[x]$  is irreducible (i.e. in  $\mathbb{Q}[x]$ )?

eg.  $f(x) = x^4 + x^2 + x + 1$

If  $f(x) = \underbrace{(x^2+ax+b)}_{\text{degree 2 in } \mathbb{Z}[x]} \underbrace{(x^2+cx+d)}_{\text{degree 2 in } \mathbb{Z}[x]}$  then  $bd=1$  implies  $b=d=\pm 1$ . If  $b=d=1$  then  $f(x) = (x^2+ax+1)(x^2-cx+1)$  has no  $x$  term, a contradiction.

If  $b=d=-1$  then  $f(x) = (x^2+ax-1)(x^2-cx-1)$  has no  $x$  term again a contradiction.

If  $f(x) = (x+a)(x^3+bx^2+cx+d)$  then  $ad=1$  so  $a=d=\pm 1$ , but  $f(1) = 4$   $\left\{ \begin{array}{l} \neq \pm 1 \text{ are not roots} \\ f(-1) = 2 \end{array} \right\}$  of  $f(x)$ .

So  $f(x)$  is irreducible in  $\mathbb{Z}[x]$ ; so  $f(x)$  is irreducible also in  $\mathbb{Q}[x]$ .

Why do we care about automorphisms of fields?

Historically the study of fields originated in questions about finding roots of polynomials.

The roots of  $ax^2+bx+c$  ( $a \neq 0$ ) are  $\frac{-b \pm \sqrt{b^2-4ac}}{2a}$ .

Similarly the roots of  $ax^3+bx^2+cx+d$  ( $a \neq 0$ ) are given explicitly using formulas of  $a, b, c, d$  using  $+, -, \times, \div$  and extracting square roots and cube roots.

Similarly for polynomials of degree 4. But for degree  $\geq 5$ , no such formula exists. The reason is found in group theory. Galois theory gives the connection between fields and groups.

Given a polynomial  $f(x) = x^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0 \in \mathbb{Q}[x]$  then  $f(x) = (x-r_1)(x-r_2)\dots(x-r_n)$  where  $r_1, \dots, r_n \in \mathbb{C}$ . The roots lie in  $F = \mathbb{Q}(r_1, \dots, r_n) \subset \mathbb{C}$ . Let  $G = \text{Aut } F$ .  $G$  permutes  $r_1, \dots, r_n$  (in particular  $G$  is a subgroup of  $S_n$ ) order  $n!$

If  $F$  is a field then  $F[a] =$  ring of all polynomials in  $a$  with all coefficients in  $F$ .

$=$  the smallest ring containing  $F$  and  $a$

$F(a) =$  the field of all rational functions in  $a$  with coefficients in  $F$

$=$  the smallest field extension of  $F$  containing  $a$ .

You can do all this for more than one element  $a$  e.g.

$F[a_1, \dots, a_k] =$  the ring of all polynomials in  $a_1, \dots, a_k$  with coefficients in  $F$

$=$  the smallest ring containing  $F$  and  $a_1, \dots, a_k$

$=$  the ring generated by  $F, a_1, \dots, a_k$

$F(a_1, \dots, a_k) =$  the field extension of  $F$  generated by  $a_1, \dots, a_k$  together with  $F$ .

eg.  $\mathbb{Q}[\sqrt{2}] = \{a_0 + a_1\sqrt{2} + a_2\sqrt{2}^2 + a_3\sqrt{2}^3 + \dots + a_n\sqrt{2}^n : n \geq 0, a_i \in \mathbb{Q}\} = \{a + b\sqrt{2} : a, b \in \mathbb{Q}\}$

$\mathbb{Q}(\sqrt{2}) = \mathbb{Q}[\sqrt{2}]$  since  $\sqrt{2}$  is algebraic.

$E = \mathbb{Q}[\sqrt{2}, \sqrt{5}]$  ... is this a field?  $\mathbb{Q}[\sqrt{2}, \sqrt{5}] = \{a + b\sqrt{2} + c\sqrt{5} + d\sqrt{10} : a, b, c, d \in \mathbb{Q}\}$

eg.  $\alpha = \sqrt{2} + \sqrt{5} \in \mathbb{Q}[\sqrt{2}, \sqrt{5}]$  is a root of a polynomial  $f(x) \in \mathbb{Q}[x]$ , in fact  $f(x) \in \mathbb{Z}[x]$ .

In fact  $\alpha \notin \mathbb{Q}$  (why?)

$$\alpha = \sqrt{2} + \sqrt{5}$$

$$\alpha^2 = 7 + 2\sqrt{10}$$

$$\alpha^2 - 7 = 2\sqrt{10}$$

$$\alpha^4 - 14\alpha^2 + 49 = 40$$

$$\alpha^4 - 14\alpha^2 + 9 = 0$$

Candidate:  $x^4 - 14x^2 + 9$

You can check that this poly. is irred. in  $\mathbb{Q}[x]$

(using steps we used on Friday Sept 13).

If  $r(x) \neq 0$  then take

$$d(x) = \gcd(f(x), r(x)) = a(x)f(x) + b(x)r(x)$$

by Euclid's Algorithm

$$d(x) = a(x)f(x) + b(x)r(x) = 0$$

Contradiction since  $f(x)$  is irreducible in  $\mathbb{Q}[x]$ .

$f(x) = x^4 - 14x^2 + 9$  is the minimal polynomial of  $\alpha$  over  $\mathbb{Q}$  in the sense that a polynomial  $g(x) \in \mathbb{Q}[x]$  has  $\alpha$  as a root iff  $f(x) \mid g(x)$  i.e.  $g(x) = u(x)f(x)$ ,  $u(x) \in \mathbb{Q}[x]$ .

Proof: If  $g(x) = u(x)f(x)$  for some  $u(x) \in \mathbb{Q}[x]$  then

$$g(\alpha) = u(\alpha)f(\alpha) = 0 \text{ i.e. } g(x) \text{ is a poly. with coeffs in } \mathbb{Q}$$

having  $\alpha$  as a root. Conversely, suppose  $g(x) \in \mathbb{Q}[x]$  having

$\alpha$  as a root. Then  $g(x) = q(x)f(x) + r(x)$  with  $q(x), r(x) \in \mathbb{Q}[x]$ ,  $\deg r(x) < 4$ .

Now  $\underbrace{g(\alpha)}_0 = \underbrace{q(\alpha)f(\alpha)}_0 + r(\alpha) = 0 \Rightarrow r(\alpha) = 0$

If  $\alpha \in \mathbb{C}$  is algebraic ( $\alpha$  is a root of coefficients in  $\mathbb{Q}$ ) then there is a minimal polynomial  $m_\alpha(x) \in \mathbb{Q}[x]$  of smallest degree which is monic i.e. its leading coeff. is 1. unique

The minimal poly. of  $\sqrt{2} + \sqrt{5}$  is  $x^4 - 14x^2 + 9 = (x^4 - 14x^2 + 49) - 40 = (x^2 - 7)^2 - 40 = (x^2 - 7 + 2\sqrt{10})(x^2 - 7 - 2\sqrt{10})$   
 The roots  $\sqrt{2} + \sqrt{5}, \sqrt{2} - \sqrt{5}, -\sqrt{2} + \sqrt{5}, -\sqrt{2} - \sqrt{5}$   
 $= (x - (\sqrt{2} + \sqrt{5}))(x - (\sqrt{2} - \sqrt{5}))(x - (-\sqrt{2} + \sqrt{5}))(x - (-\sqrt{2} - \sqrt{5}))$   
 $= (x + \sqrt{2} - \sqrt{5})(x - \sqrt{2} + \sqrt{5})(x - \sqrt{2} - \sqrt{5})(x + \sqrt{2} + \sqrt{5})$

$$\sqrt{7 - 2\sqrt{10}} = -\sqrt{2} + \sqrt{5} \quad \text{since } (-\sqrt{2} + \sqrt{5})^2 = 7 - 2\sqrt{10}$$

$$(\sqrt{2} - \sqrt{5})^2 = 7 - 2\sqrt{10}$$

$$\sqrt{7 + 2\sqrt{10}} = \sqrt{2} + \sqrt{5} \quad \text{since } (\sqrt{2} + \sqrt{5})^2 = 7 + 2\sqrt{10}$$

$$(-\sqrt{2} - \sqrt{5})^2 = 7 + 2\sqrt{10}$$

$\sqrt{2} \notin \mathbb{Q}$  by Euclid's argument

If  $\sqrt{2} = \frac{m}{n}$ ,  $m, n \in \mathbb{Z}$  in lowest terms i.e.  $\gcd(m, n) = 1$   
 then  $m^2 = 2n^2$  is even so  $m = 2r$ ,  $r \in \mathbb{Z}$ ,  $4r^2 = 2n^2$ ,  $n^2 = 2r^2$   
 is even so  $n$  is even, a contradiction.

The same argument shows  $\sqrt{5}, \sqrt{10} \notin \mathbb{Q}$ .

$\pm\sqrt{2} \pm \sqrt{5} \notin \mathbb{Q}$ , since their squares are  $7 \pm 2\sqrt{10} \notin \mathbb{Q}$ .  
 This gives another explanation why  $x^4 - 14x^2 + 9$  is irreducible in  $\mathbb{Q}[x]$ .

$$E = \mathbb{Q}[\sqrt{2}, \sqrt{5}] = \mathbb{Q}[\alpha], \quad \alpha = \sqrt{2} + \sqrt{5}$$

$$\{a + b\sqrt{2} + c\sqrt{5} + d\sqrt{10} : a, b, c, d \in \mathbb{Q}\} = \{a + b\alpha + c\alpha^2 + d\alpha^3 : a, b, c, d \in \mathbb{Q}\}$$

This equality is explained as follows:  $E = \mathbb{Q}[\sqrt{2}, \sqrt{5}] = \{a + b\sqrt{2} + c\sqrt{5} + d\sqrt{10} : a, b, c, d \in \mathbb{Q}\}$   
 is a 4-dimensional vector space over  $\mathbb{Q}$  with basis  $\{1, \sqrt{2}, \sqrt{5}, \sqrt{10}\}$ .

$$E \supseteq \mathbb{Q}[\alpha], \quad \alpha^4 - 14\alpha^2 + 9 = 0 \quad \text{so}$$

$$\alpha^4 = 14\alpha^2 - 9$$

$$\alpha^5 = 14\alpha^3 - 9\alpha$$

$$\alpha^6 = 14\alpha^4 - 9\alpha^2 = 14(14\alpha^2 - 9) - 9\alpha^2 = 187\alpha^2 - 126$$

$$\{a + b\alpha + c\alpha^2 + d\alpha^3 : a, b, c, d \in \mathbb{Q}\}$$

$$\{a, b, c, d \in \mathbb{Q}\}$$

There is no nonzero  $(a, b, c, d) \in \mathbb{Q}^4$  with  $a + b\alpha + c\alpha^2 + d\alpha^3 = 0$



So  $1, \alpha, \alpha^2, \alpha^3$  are linearly independent over  $\mathbb{Q}$ .

An important class of examples of fields is: (algebraic) number fields are finite-dimensional extensions  $E \supseteq \mathbb{Q}$  eg.

$\mathbb{Q}, \mathbb{Q}[\sqrt{2}], \mathbb{Q}[\sqrt{5}], \mathbb{Q}[i], \mathbb{Q}[\sqrt{-3}], \mathbb{Q}[\alpha] = \mathbb{Q}[\sqrt{2}, \sqrt{5}]$ , etc.  
 $\alpha = \sqrt{2} + \sqrt{5}$

$\langle \sigma \rangle = \{1, \sigma\}$   
 $\langle \tau \rangle = \{1, \tau\}$   
 $\langle \sigma\tau \rangle = \{1, \sigma\tau\}$   
 $\langle 1 \rangle = \{1\}$

Not  $\mathbb{R}, \mathbb{C}$  which are infinite-dimensional over  $\mathbb{Q}$ .  
 $1, \sqrt{2}, \sqrt{3}, \sqrt{5}, \sqrt{7}, \sqrt{11}, \sqrt{13}, \dots$  are linearly independent over  $\mathbb{Q}$ .



There is a one-to-one correspondence between subgroups of  $G = \text{Aut } E$  and subfields of  $E$ .  
 Corresponding to every subfield  $K \subseteq E$ , we have the subgroup  $G_K = \{\phi \in \text{Aut } E : \phi \text{ fixes every element of } K\}$ .

This diagram is a Hasse diagram showing all subfields of  $E$ .

This fact is not quite obvious.

$\text{Aut } E = \{1, \sigma, \tau, \sigma\tau\}$   
 identity

$\sigma^2 = 1$   
 $\tau^2 = 1$

$\phi(\sqrt{2})\phi(\sqrt{2}) = \phi(\sqrt{2} \cdot \sqrt{2}) = 2 \Rightarrow$

$\phi(\sqrt{2}) = \pm\sqrt{2}$   
 $\phi(\sqrt{5}) = \pm\sqrt{5}$   
 $\phi(\sqrt{10}) = \pm\sqrt{10}$

$\sigma(\sqrt{10}) = \sigma(\sqrt{2}\sqrt{5}) = \sigma(\sqrt{2})\sigma(\sqrt{5}) = \sqrt{2}(-\sqrt{5}) = -\sqrt{10}$

$\begin{matrix} H \\ \downarrow \\ L \end{matrix}$

$k = [H:L] = \text{index of } L \text{ in } H = \text{number of cosets of } L \text{ in } H = \frac{|H|}{|L|}$

$a$	1	$\sqrt{2}$	$\sqrt{5}$	$\sqrt{10}$
$1(a)$	1	$\sqrt{2}$	$\sqrt{5}$	$\sqrt{10}$
$\sigma(a)$	1	$\sqrt{2}$	$-\sqrt{5}$	$-\sqrt{10}$
$\tau(a)$	1	$-\sqrt{2}$	$\sqrt{5}$	$-\sqrt{10}$
$\sigma\tau(a)$	1	$-\sqrt{2}$	$-\sqrt{5}$	$\sqrt{10}$

Aut  $E$  is the Klein four-group.

Every automorphism  $\phi: E \rightarrow E$  satisfies  $\phi(a) = a$  for all  $a \in \mathbb{Q}$ .  
 So  $\phi$  is determined by  $\phi(\sqrt{2}), \phi(\sqrt{5}), \phi(\sqrt{10})$ .

$\phi(a + b\sqrt{2} + c\sqrt{5} + d\sqrt{10}) = \phi(a) + \phi(b)\phi(\sqrt{2}) + \phi(c)\phi(\sqrt{5}) + \phi(d)\phi(\sqrt{10})$   
 $= a + b\phi(\sqrt{2}) + c\phi(\sqrt{5}) + d\phi(\sqrt{10})$   
 $a, b, c, d \in \mathbb{Q}$



If  $F \subseteq E$  is a subfield ( $F$  is a subfield of  $\cdot$ , i.e.  $E$  is an extension of  $F$ ) then  $E$  is a vector space over  $F$ . The dimension of  $E$  over  $F$  is the degree of  $E$  over  $F$ , denoted  $[E:F]$ .

eg.  $[\mathbb{C}:\mathbb{R}] = 2$  with basis  $\{1, i\}$ .


$[\mathbb{Q}[\sqrt{2}):\mathbb{Q}] = 2$  with basis  $\{1, \sqrt{2}\}$ .

$[\mathbb{Q}[\sqrt{2}, \sqrt{5}):\mathbb{Q}] = 4$  with basis  $\{1, \sqrt{2}, \sqrt{5}, \sqrt{10}\}$  or  $\{1, \alpha, \alpha^2, \alpha^3\}$  where  $\alpha = \sqrt{2} + \sqrt{5}$

$[E:\mathbb{Q}[\sqrt{2}]] = 2$  with basis  $\{1, \sqrt{5}\}$

Elements of  $E$  have the form

$$a + b\sqrt{2} + c\sqrt{5} + d\sqrt{10} = (a + b\sqrt{2}) + (c + d\sqrt{2})\sqrt{5}$$



Theorem If  $F \subseteq K \subseteq E$  is a tower of fields (i.e. subfields/extensions) then  $[E:F] = [E:K][K:F]$ .

In fact if  $[K:F] = m$  with basis  $\{\alpha_1, \dots, \alpha_m\}$  for  $K$  over  $F$  and  $[E:K] = n$  with basis  $\{\beta_1, \dots, \beta_n\}$  for  $E$  over  $K$  then  $\{\alpha_i \beta_j : 1 \leq i \leq m, 1 \leq j \leq n\}$  is a basis for  $E$  over  $F$ .

One of these is in  $\mathbb{Q}[\sqrt{2}]$ :

$\sqrt{3+2\sqrt{2}} = 1 + \sqrt{2} \in \mathbb{Q}[\sqrt{2}]$  since  $(1+\sqrt{2})^2 = 1+2+2\sqrt{2} = 3+2\sqrt{2}$

let  $\alpha = \sqrt{2+3\sqrt{2}} \notin \mathbb{Q}[\sqrt{2}]$ . (but this is not yet clear)

Let  $\alpha = \sqrt{2+3\sqrt{2}}$ . Then  $\alpha$  is algebraic. Find the minimal poly. of  $\alpha$  over  $\mathbb{Q}$  (the unique minic poly. of smallest degree in  $\mathbb{Q}[x]$  having  $\alpha$  as a root).

$$\begin{aligned}\alpha &= \sqrt{2+3\sqrt{2}} \\ \alpha^2 &= 2+3\sqrt{2} \\ \alpha^2 - 2 &= 3\sqrt{2} \\ \alpha^4 - 4\alpha^2 + 4 &= 18 \\ \alpha^4 - 4\alpha^2 - 14 &= 0\end{aligned}$$

$f(x) = x^4 - 4x^2 - 14$  is the minimal poly. of  $\alpha$  over  $\mathbb{Q}$ .  
To see this, we need to check that  $f(x)$  is irreducible in  $\mathbb{Q}[x]$  (equivalently, in  $\mathbb{Z}[x]$ ).

The four roots of  $f(x)$  in  $\mathbb{C}$  are

$$\begin{aligned}\alpha &= \sqrt{2+3\sqrt{2}} \\ -\alpha &= -\sqrt{2+3\sqrt{2}} \\ \beta &= \sqrt{2-3\sqrt{2}} \\ -\beta &= -\sqrt{2-3\sqrt{2}}\end{aligned}$$

$$\begin{aligned}\beta &= \sqrt{2-3\sqrt{2}} \\ \beta^2 &= 2-3\sqrt{2} \\ \beta^2 - 2 &= -3\sqrt{2} \\ \beta^4 - 4\beta^2 + 4 &= 18 \\ \beta^4 - 4\beta^2 - 14 &= 0\end{aligned}$$

$$\begin{aligned}\text{So } f(x) &= x^4 - 4x^2 - 14 \\ &= (x-\alpha)(x+\alpha)(x-\beta)(x+\beta) \text{ in } \mathbb{C}[x]\end{aligned}$$

where  $(x-\alpha)(x+\alpha) = x^2 - \alpha^2 = x^2 - (2+3\sqrt{2}) \notin \mathbb{Q}[x]$

$(x-\alpha)(x-\beta) \notin \mathbb{Q}[x]$ .

$\alpha \in \mathbb{Q}[\alpha] = \{a + b\alpha + c\alpha^2 + d\alpha^3 : a, b, c, d \in \mathbb{Q}\}$   
has  $\{1, \alpha, \alpha^2, \alpha^3\}$  as a basis over  $\mathbb{Q}$ .

$$[\mathbb{Q}[\alpha] : \mathbb{Q}] = 4$$

$\mathbb{Q}[\alpha]$

$\supseteq$

$\mathbb{Q}[\sqrt{2}]$

$\supseteq$

$\mathbb{Q}$

$\xleftrightarrow{\text{no Galois correspondence}} \langle \sigma \rangle = \text{Aut } E$

$\supseteq$

$$\alpha^2 = 2+3\sqrt{2} \Rightarrow \sqrt{2} = \frac{1}{3}\alpha^2 - \frac{2}{3} \in \mathbb{Q}[\alpha]$$

$$\alpha^4 = 4\alpha^2 + 14$$

$$\alpha^5 = 4\alpha^3 + 14\alpha^2$$

$$\alpha^6 = 4\alpha^4 + 14\alpha^3 = 4(4\alpha^2 + 14) + 14\alpha^3 = 14\alpha^3 + 16\alpha^2 + 56$$

$$\begin{aligned}\sigma(\alpha^3) &= \sigma(\alpha)\sigma(\alpha)\sigma(\alpha) \\ &= \sigma(\alpha)\sigma(\alpha^2) \\ &= \sigma(\alpha)^3\end{aligned}$$

What are the automorphisms of  $E = \mathbb{Q}[\alpha]$ ?  $\iota = \text{identity}$ ,  $\iota(x) = x$ , is an automorphism.  
A function  $\sigma$  mapping  $\alpha \mapsto -\alpha$  is an automorphism.

$$\begin{aligned}\text{Note: } \sigma(a + b\alpha + c\alpha^2 + d\alpha^3) &= \sigma(a) + \sigma(b\alpha) + \sigma(c\alpha^2) + \sigma(d\alpha^3) = \sigma(a) + \sigma(b)\sigma(\alpha) + \sigma(c)\sigma(\alpha^2) + \sigma(d)\sigma(\alpha^3) \\ &= a - b\alpha + c\alpha^2 - d\alpha^3\end{aligned}$$

$(a, b, c, d \in \mathbb{Q})$

Note:  $\sigma^2 = \iota$

$$\alpha = \sqrt{2+3\sqrt{2}}$$

$$-\alpha = -\sqrt{2+3\sqrt{2}}$$

$$\beta = \sqrt{2-3\sqrt{2}}$$

$$-\beta = -\sqrt{2-3\sqrt{2}}$$

There is also an automorphism  $\phi: \alpha \mapsto \beta$ . (There is only one such automorphism.)

$$\phi(a + b\alpha + c\alpha^2 + d\alpha^3) = a + b\beta + c\beta^2 + d\beta^3 \quad \text{for all } a, b, c, d \in \mathbb{Q}.$$

This is  $\phi(x)$ ,  $x \in E = \mathbb{Q}[\alpha]$ . What is  $\phi^2(x)$ ?

$$\phi^2(a + b\alpha + c\alpha^2 + d\alpha^3) = ?$$

$$\phi^2(\alpha) = \phi(\phi(\alpha)) = \phi(\beta) = ?$$

First write  $\beta$  in the standard form  $(*) + (*)\alpha + (*)\alpha^2 + (*)\alpha^3$ . This is impossible.

Any automorphism of  $E = \mathbb{Q}[\alpha]$  must map  $\alpha$  to a root of  $f(x) = x^4 - 4x^2 - 14$ . Why?

$$\alpha^4 - 4\alpha^2 - 14 = 0$$

Apply  $\phi \in \text{Aut } E$  to both sides.

$$\phi(\alpha^4 - 4\alpha^2 - 14) = \phi(0) = 0$$

$$\phi(\alpha^4) - 4\phi(\alpha^2) - \phi(14) = 0$$

$$\phi(\alpha)^4 - 4\phi(\alpha)^2 - 14 = 0$$

$$f(\phi(\alpha)) = 0 \Rightarrow \phi(\alpha) \in \{\pm\alpha, \pm\beta\}.$$

But if  $\phi(\alpha) = \beta$  then  $\phi \notin \text{Aut } E$ .

$$\begin{array}{c} \uparrow \quad \quad \uparrow \\ \alpha \in \mathbb{R} \quad \beta \notin \mathbb{R} \end{array}$$

Actually  $E = \mathbb{Q}[\alpha]$  has only two automorphisms 1,  $\phi$ .

The extension  $E \supset \mathbb{Q}$  does not contain all the roots of  $f(x) = x^4 - 4x^2 - 14$ .

An extension  $F[\alpha] \supseteq F$  is normal if  $F[\alpha]$  contains all the roots of the min. poly.  $f(x)$  of  $\alpha$  over  $F$ .

$$\text{Eg. } \alpha = \sqrt{7+\sqrt{2}}$$

$$\alpha^2 = 7+\sqrt{2}$$

$$\alpha^2 - 7 = \sqrt{2}$$

$$\alpha^4 - 14\alpha^2 + 49 = 2$$

$$\alpha^4 - 14\alpha^2 + 47 = 0$$

The minimal poly. of  $\alpha$  over  $\mathbb{Q}$  is  $f(x) = x^4 - 14x^2 + 47 \in \mathbb{Q}[x]$ .

(Exercise:  $f(x)$  is irreducible in  $\mathbb{Q}[x]$  so it really is the min. poly. of  $\alpha$  over  $\mathbb{Q}$ .)

The roots of  $f(x)$  are

$$\alpha = \sqrt{7+\sqrt{2}}$$

$$-\alpha = -\sqrt{7+\sqrt{2}}$$

$$\beta = \sqrt{7-\sqrt{2}}$$

$$-\beta = -\sqrt{7-\sqrt{2}}$$

$$f(x) = x^4 - 14x^2 + 47 = (x-\alpha)(x+\alpha)(x-\beta)(x+\beta)$$

In this case  $E = \mathbb{Q}[\alpha] = \{a + b\alpha + c\alpha^2 + d\alpha^3 : a, b, c, d \in \mathbb{Q}\}$  contains all the roots of  $f(x)$

so it is a normal extension of  $\mathbb{Q}$ .

$$\beta = (*) + (*)\alpha + (*)\alpha^2 + (*)\alpha^3$$