



# Field Theory

Book 2

Claim:  $\sqrt{3} \notin \mathbb{Q}[\sqrt{2}]$  i.e.  $\sqrt{3} = a + b\sqrt{2}$  has no solution with  $a, b \in \mathbb{Q}$ .

Suppose  $\sqrt{3} = a + b\sqrt{2}$  where  $a, b \in \mathbb{Q}$ . Then  $3 = a^2 + 2b^2 + 2ab\sqrt{2}$  so  $2ab\sqrt{2} = 3 - a^2 - 2b^2$ .

If  $ab \neq 0$  then  $\sqrt{2} = \frac{3 - a^2 - 2b^2}{2ab} \in \mathbb{Q}$ , a contradiction.

If  $b = 0$  then  $0 = 3 - a^2$  so  $a^2 = 3$ ,  $a = \pm\sqrt{3} \notin \mathbb{Q}$ , a contradiction.

If  $a = 0$  then  $0 = 3 - 2b^2$  so  $2b^2 = 3$ ,  $4b^2 = 6$ ,  $2b = \pm\sqrt{6} \notin \mathbb{Q}$ , a contradiction.  $\square$

So  $1, \sqrt{2}, \sqrt{3} \in \mathbb{R}$  are linearly independent over  $\mathbb{Q}$ .

•  $1 \neq 0$

•  $\sqrt{2} \neq$  scalar multiple of  $1$ . ( $\sqrt{2} \notin \mathbb{Q}$  by Euclid)

•  $\sqrt{3} \neq$  linear combination of  $1, \sqrt{2}$ . (proved above)

$$\sqrt{8} = 2\sqrt{2}$$

$1, \sqrt{2}, \sqrt{3}, \sqrt{5}, \sqrt{7}, \sqrt{11}, \sqrt{13}, \sqrt{17}, \sqrt{19}, \sqrt{23}, \dots$  are linearly independent.

$[\mathbb{R} : \mathbb{Q}] = \infty$  (in fact uncountable)

Also  $1, \pi, \pi^2, \pi^3, \dots$  are linearly independent over  $\mathbb{Q}$ . (since  $\pi$  is transcendental).

An extension  $E \supseteq F$  is finite if  $[E : F] < \infty$ , i.e.  $[E : F] = n$  is a positive integer.

eg  $\mathbb{C} \supseteq \mathbb{R}$  is a quadratic extension, hence finite,  $[\mathbb{C} : \mathbb{R}] = 2$ .

A finite extension of  $\mathbb{Q}$  i.e.  $E \supseteq \mathbb{Q}$  with  $[E : \mathbb{Q}] = n$ , a positive integer, is called a number field (or algebraic number field). Here every element  $\alpha \in E$  is algebraic over  $\mathbb{Q}$ . Why?

$1, \alpha, \alpha^2, \dots, \alpha^n$  are  $n+1$  vectors in an  $n$ -dimensional vector space  $E \supseteq \mathbb{Q}$  so this list is linearly dependent i.e.  $a_0 + a_1\alpha + a_2\alpha^2 + \dots + a_n\alpha^n = 0$  for some  $a_0, a_1, \dots, a_n \in \mathbb{Q}$ , not all zero, i.e.  $\alpha$  is a root of some nonzero polynomial  $f(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n \in \mathbb{Q}[x]$ .

If  $f(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$  then the degree of  $f(x)$ , denoted  $\deg f(x)$ , is the largest  $d$  such that  $a_d \neq 0$ .

$$\deg(3x^2 + 5x + 7) = 2$$

$$\deg(0x^2 + 5x + 7) = \deg(5x + 7) = 1$$

$$\deg(7) = \deg(7x^0) = 0$$

$\deg 0$  is sometimes left undefined (not 0) or  $\deg 0 = -\infty$ .

$$\deg[(3x^2 + 5x + 7)(x^3 - 4x - 11)] = \deg(3x^5 + \dots - 77) = 5$$

$$\deg[f(x)g(x)] = \deg f(x) + \deg g(x)$$

If  $g(x) = x^3 - 4x - 11$  then  $\deg 0 = 0$

$$\deg(0g(x)) = \deg 0$$

$$\deg 0 + \deg g(x) = \deg 0 + 3$$

$$\deg 0 = (\deg 0) + 3$$

There is no integer value for  $\deg 0$  that satisfies this.  
(We don't choose  $+\infty$ ; we choose  $-\infty$ .)

Let  $\alpha \in \mathbb{C}$ . If  $\alpha$  is the root of some nonzero poly.  $f(x) \in \mathbb{Q}[x]$  (i.e.  $f(\alpha) = 0$ ) then  $\alpha$  is algebraic of degree  $n$  where  $n$  is the smallest degree of any such polynomial  $f(x)$ . In this case, the smallest degree monic polynomial having  $\alpha$  as a root is the minimal polynomial of  $\alpha$  (over  $\mathbb{Q}$ ).

eg.  $\sqrt{14}$  is algebraic of degree 2 with min. poly.  $x^2 - 14 \in \mathbb{Q}[x]$ .

Look at powers  $1, \alpha, \alpha^2, \alpha^3, \dots$

$$\alpha = \sqrt{14} \Rightarrow 1, \alpha \text{ lin. indep.}$$

$$1, \alpha, \alpha^2 \text{ lin. dep.}$$

$$\alpha^2 = 0 \cdot \alpha + 14 \cdot 1$$

$\alpha = \sqrt{2} + \sqrt{5}$  is algebraic of degree 4 with min. poly.  $x^4 - 10x^2 + 1$ . Why is  $\alpha = \sqrt{2} + \sqrt{5}$  not a root of any smaller degree poly. with rational coefficients?

If  $x^4 - 10x^2 + 1 = f(x)g(x)$  where  $f(x), g(x) \in \mathbb{Q}[x]$  then one of  $f(x), g(x)$  is a <sup>(non-zero)</sup> constant polynomial. Assuming we start with a monic poly. with integer coefficients, it suffices to check that there is no nontrivial factorization over  $\mathbb{Z}[x]$ .

If  $x^4 - 10x^2 + 1 = f(x)g(x)$ ,  $f(x), g(x) \in \mathbb{Z}[x]$ , neither  $f(x)$  nor  $g(x)$  is constant then either

(i)  $\deg f(x) = 1$ ,  $\deg g(x) = 3$ ; or

(ii)  $\deg f(x) = \deg g(x) = 2$ .

(The case  $\deg f(x) = 3$ ,  $\deg g(x) = 1$  is essentially case (i)).

In both cases we obtain a contradiction.

In case (i),  $x^4 - 10x^2 + 1 = (x+a)(x^3+bx^2+cx+d)$ ,  $a=1$ ,  $a=d = \pm 1$ .