





Cyclotomic Fields

Book 1

A (complex) n^{th} root of unity is a value $\xi \in \mathbb{C}$ satisfying $\xi^n = 1$.

$$\xi = e^{2\pi i k/n}$$



Solutions of $z^n = 1$ form a ^{multiplicative} cyclic group of order n .

There are $\phi(n)$ primitive n^{th} roots of unity.

Euler's totient function

$\phi(n)$ = number of generators in any cyclic group of order n .

eg. $n=6$ $\phi(6) = 2$.


$$\phi(n) = |\{k \in \{1, 2, \dots, n\} : \gcd(k, n) = 1\}|$$

1, 5 relatively prime to 6.

All sixth roots of unity are ξ^k , $1 \leq k \leq n$. $-1 = \xi^3$

If $z^n = 1$ and $w^m = 1$ then $(zw)^{mn} = 1$

The product of an m^{th} root of unity with an n^{th} root of unity is an $(mn)^{\text{th}}$ root of unity (in fact an $\text{lcm}(m, n)^{\text{th}}$ root of unity).



A cyclotomic field is a field $\mathbb{Q}[\xi]$ where $\xi \in \mathbb{C}$ is a root of unity.

$$\xi^n = 1 \Rightarrow 1 + \xi + \xi^2 + \dots + \xi^{n-1} = 0$$

$$z^n - 1 = (z-1)(1 + z + z^2 + \dots + z^{n-1})$$

$$n=6: z^6 - 1 = \Phi_1(z) \Phi_2(z) \Phi_3(z) \Phi_6(z) \quad (\text{factorization into irreducible factors in } \mathbb{Q}[x] \text{ or } \mathbb{Z}[x])$$

Roots: $1, -1, \xi^2, \xi^4, \xi, \xi^5$
 primitive sixth roots of 1.

$\Phi_n(z)$ = minimal poly. of $\xi = e^{2\pi i/n}$ over $\mathbb{Q}$ (or over $\mathbb{Z}$)
 (simplest monic poly. with rational coefficients having ξ as a root)

$$z^n - 1 = \prod_{\substack{d|n \\ d \geq 1}} \Phi_d(z) \quad \text{product over all positive integer divisors of } n.$$

Divisors of 6 are $\pm 1, \pm 2, \pm 3, \pm 6$

Positive divisors of 6 are 1, 2, 3, 6.

$$\Phi_n(z) = \prod_{\substack{1 \leq k \leq n \\ \gcd(k, n) = 1}} (z - \xi^k) \quad \text{has as its roots the primitive } n^{\text{th}} \text{ roots of } 1.$$

$$\mathbb{Q}[\xi_n] = \{ \text{polynomials in } \xi \text{ with coefficients in } \mathbb{Q} \}$$

$$= \{ \text{linear combinations of } 1, \xi, \xi^2, \dots \text{ with rational coefficients} \}$$

$$= \{ \text{linear combinations of } 1, \xi, \xi^2, \dots, \xi^{n-1} \text{ with coefficients in } \mathbb{Q} \}$$

$$\xi = \xi_n = e^{2\pi i/n}$$

The roots of unity in $\mathbb{C}$ form a multiplicative group which is abelian but not cyclic.
 This multiplicative group is isomorphic to the additive group $\mathbb{Q}/\mathbb{Z}$.

$$\{\text{roots of unity in } \mathbb{C}\} = \{e^{i\alpha \cdot 2\pi} : \alpha \in \mathbb{Q}\} \quad \alpha \mapsto \alpha+1 \text{ doesn't change the value of } e^{i\alpha \cdot 2\pi}$$

The homomorphism $\mathbb{Q} \xrightarrow{f} \{\text{roots of unity}\}$ is onto but not one-to-one.
 $\alpha \mapsto e^{i\alpha \cdot 2\pi}$

By the first isomorphism theorem for groups, $\{\text{roots of unity}\} \cong \mathbb{Q}/\mathbb{Z}$.
 Integer linear combinations of roots of unity are cyclotomic integers. $\leftarrow \ker f$
 This is a ring.

$\mathbb{Q}$ -linear combinations of roots of unity form a field, the field of cyclotomic numbers.
 Both these (ring and field) are countably infinite.

$i = \sqrt{-1}$ is a cyclotomic integer (a 4th root of unity)

$\sqrt{7}$ is a cyclotomic integer. (not really obvious)

$\sqrt[3]{7} = 7^{1/3}$ is not

$\xi = \xi_7 = e^{2\pi i/7}$ is a primitive 7th root of unity

$\alpha = \xi + \xi^2 + \xi^4$ is a cyclotomic integer



$$\alpha^2 = \xi^2 + \xi^4 + \xi + 2\xi^3 + 2\xi^5 + 2\xi^6$$

$$0 = 2 + 2\xi + 2\xi^2 + 2\xi^3 + 2\xi^4 + 2\xi^5 + 2\xi^6$$

$$\alpha^2 = -2 - \xi - \xi^2 - \xi^4 = 2 - \alpha$$

$$\alpha^2 + \alpha + 2 = 0 \Rightarrow \alpha = \frac{-1 \pm \sqrt{1-8}}{2} = \frac{-1 \pm \sqrt{-7}}{2} = \frac{-1 \pm i\sqrt{7}}{2}$$

$$2\alpha + 1 = \pm i\sqrt{7}$$



$$2\alpha + 1 = \pm i\sqrt{7}$$

$$\mp i(2\alpha + 1) = \sqrt{7}$$

$$\alpha = \xi + \xi^2 + \xi^4$$

$$\mp i(2\alpha + 1) = \mp i(1 + 2e^{2\pi i/7} + 2e^{4\pi i/7} + 2e^{8\pi i/7}) = \sqrt{7}$$

2.645751311... $\swarrow$

$\searrow$ cosine terms

$$= \pm \left(2\sin\frac{2\pi}{7} + 2\sin\frac{4\pi}{7} + \sin\frac{8\pi}{7} \right) + i(\dots)$$

$$2.645751311\dots$$

$$\sqrt{7} = 2\sin\frac{2\pi}{7} + 2\sin\frac{4\pi}{7} + 2\sin\frac{8\pi}{7} \text{ is a cyclotomic integer.}$$

$$2\sin\frac{2\pi}{7} = -i(\xi - \xi^6) = -i\left[\left(\cos\frac{2\pi}{7} + i\sin\frac{2\pi}{7}\right) - \left(\cos\frac{2\pi}{7} - i\sin\frac{2\pi}{7}\right)\right]$$