




Math 5605

Algebraic Topology


Book 2

When are two covering maps of X equivalent? Say $Y \xrightarrow{f} X$, $Y' \xrightarrow{f'} X$ are covering maps.





Graph i.e. combinatorial graph with vertices $\{1, 2, 3, 4\}$ and edges $\{1, 2\}, \{1, 3\}, \dots, \{3, 4\}$.

X is the geometric realization of this graph formed as a disjoint union of copies of $[0, 1]$ with endpoints identified as required by the picture.





A homomorphism of graphs $\Gamma \xrightarrow{f} \Gamma'$ is a map $V(\Gamma) \xrightarrow{f} V(\Gamma')$ preserving adjacency i.e. $x \sim y$ in $\Gamma \Rightarrow f(x) \sim f(y)$ in Γ' . A covering map of graphs is a homomorphism $(x, y \in V(\Gamma), \{x, y\} \in E(\Gamma))$ inducing a bijection on the neighbours of each vertex of Γ (and the preimage of the neighbours of each vertex $y \in \Gamma'$ are copies of the neighbours of y).




The covering space is $Y \rightarrow X$ (or informally just Y). Some other double covers of X :

Trivial double cover




When are two covers of X equivalent (isomorphic, i.e. essentially the same)?


Let $p_1: X_1 \rightarrow X$, $p_2: X_2 \rightarrow X$ be covering spaces of X . We say $\theta: X_1 \rightarrow X_2$ is an equivalence or isomorphism of the two covers if θ is a homeomorphism and $p_2 \circ \theta = p_1$, i.e.




Ex.






is not equivalent to $Y \rightarrow X$. But what about




Is this equivalent to $Z \rightarrow X$? No...




Another picture of these covers:



$X \sqcup X$





Y




W

To construct an r -fold cover of X , create one copy of $[r] = \{1, 2, \dots, r\}$ for each vertex of X . Then for each edge of X , match up the corresponding fibres in the cover using a chosen permutation.



A triple cover $Y \rightarrow X$ is constructed as



Why is 2 more special than other positive integers (the oddest prime of all)?




has many triple covers including



The covering maps $Y_1 \rightarrow X$ and $Y_2 \rightarrow X$ are not equivalent.

An equivalence between $Y \rightarrow X$ and itself (automorphism of the cover) is a deck transformation. This is the same as a homeomorphism $Y \rightarrow Y$ which preserves fibres.

In the example above, $Y_2 \rightarrow X$ has 3 automorphisms (deck transformations) but $Y_1 \rightarrow X$ has only one (trivial) deck transformation.



In a connected r -fold cover, there are at most r deck transformations. If equality holds, the covering space is normal or Galois. (not the same as normal space in point set topology). Double covers are always normal.






In group theory, subgroups of index 2 are normal.

In the case of ^(separable) extensions of fields, the extension is normal.

For a field extension $E \supseteq F$, the degree of the extension is $[E:F]$ = dimension of E as a vector space over F . The number of F -automorphisms of E (i.e. $\sigma: E \rightarrow E$ automorphism fixing $\sigma(a) = a$ for all $a \in F$) is at most $[E:F]$. If this number is equal, it's a normal or Galois extension. Extensions of degree 2 (quadratic extensions) are always normal.

Double covers: examples



S^n is not a top. group unless $n \in \{1, 3\}$.

$$S^1 = \{z \in \mathbb{C} : |z| = 1\}$$


$$S^3 = \{z \in \mathbb{H} : |z| = 1\} \quad \mathbb{H} = \{a+bi+cj+dk : a, b, c, d \in \mathbb{R}\} \quad i^2 = j^2 = k^2 = ijk = -1$$

$$\cong SU_2(\mathbb{C}) = \left\{ A = \begin{bmatrix} \alpha & \beta \\ \gamma & \delta \end{bmatrix} : \alpha, \beta, \gamma, \delta \in \mathbb{C}, AA^* = A^*A = I, \det A = 1 \right\}$$

$$SO_3(\mathbb{R}) = \{A \in \mathbb{R}^{3 \times 3} : AA^T = A^T A = I, \det A = 1\}$$


$$O_3(\mathbb{R}) = \{A \in \mathbb{R}^{3 \times 3} : AA^T = A^T A = I\} \text{ has two connected components}$$

Fact: $S^3 \cong SU_2(\mathbb{C}) \rightarrow SO_3(\mathbb{R})$ is a double cover. $Z(S^3) = \{\pm 1\}$ homeomorphism
 $PSU_2(\mathbb{C}) = S^3 / Z(S^3) \cong SO_3(\mathbb{R}) \cong \mathbb{P}^3$




In general for $n \geq 3$, $\pi_1(SO_n(\mathbb{R})) \cong \mathbb{Z}/2\mathbb{Z}$.

$Spin_n(\mathbb{R}) \longrightarrow SO_n(\mathbb{R})$ is its universal cover; a double cover constructed from Clifford Algebras (generalizing #1). simply connected



$Y = T^2$



$f_0: [0,1] \rightarrow X$

$f_1: [0,1] \rightarrow X$

Assuming X is path-connected and $p: Y \rightarrow X$ is a path-connected covering space,


$X = Y/\sim$ where two points $y_0, y_1 \in Y$ satisfy $y_0 \sim y_1$ iff

$p(y_0) = p(y_1)$.


$f_0: [0,1] \rightarrow X$
path in X
from $f_0(0) = x_0$
to $f_0(1) = x_1$

$f_1: [0,1] \rightarrow X$
is another path in X
from x_0 to x_1
homotopic to f_0

In any covering space $p: Y \rightarrow X$ and given any path $f: [0,1] \rightarrow X$ starting at $f(0) = x_0$, the path f can be lifted to Y i.e. there is a path $g: [0,1] \rightarrow Y$ such that $f = p \circ g$ i.e.



and this lift is unique if we say which of the points in $f(x)$ to take as the starting point for g .



$$p^{-1}(x_0) = \{y_0, y_1, y_2, \dots\}$$


$$p^{-1}(x_1) = \{z_0, z_1, z_2, \dots\}$$

Every path f in X from x_0 to x_1 gives a bijection between fibres $p^{-1}(x_0) \rightarrow p^{-1}(x_1)$.

In particular if p is k -to-1 at x_0 i.e.


$|p^{-1}(x_0)| = k$ then it is k -to-1 everywhere i.e. $|p^{-1}(x)| = k$ for all $x \in X$.

More generally, if f_t is a homotopy in X and we are given f_0 , then every lifting of f_0 to Y extends to a lifting of f_t to Y .



\mathbb{R}^2 is the universal cover of T^2

$$\mathbb{R}^2 \rightarrow T^2 = \mathbb{R}^2 / \mathbb{Z}^2$$




Which top. spaces have a universal cover? (equivalently a simply connected cover)

Let X be a path-connected space. Then X has a path-connected and universal cover iff X is

- path-connected
- locally path-connected
- semi-locally simply connected



Example of a top. space without a universal cover: Hawaiian earring $\subset \mathbb{R}^2$



(not a CW complex)

$$\neq \underbrace{S^1 \vee S^1 \vee S^1 \vee \dots}_{\text{countable wedge sum (CW complex)}}$$


countable wedge sum (CW complex)



Universal cover of K_4




trivalent tree



(also the universal cover of any trivalent connected graph)
i.e. regular of degree 3 connected

Universal cover of any connected regular graph of degree $\neq 1$ is




Cayley graph of $\text{Free}\{a,b\} = G$
 Vertices correspond to elements of G .


Every vertex $w \in G$ has
 edges to wa, wa^{-1}, wb, wb^{-1} .



Universal cover of $K_{3,4}$



$$\tilde{X} = X/G$$



$\mathbb{P}^2\mathbb{R}$ has S^2 as its universal cover.

$$S^2 \longrightarrow \mathbb{P}^2\mathbb{R} = S^2/G$$

quotient of S^2
 by the antipodal
 relation.

$G = \{1, -1\}$ acts on S^2

$$1x = x$$

$$(-1)x = -x \quad (\text{antipode of } x)$$

X/\sim = partition of X into equivalence classes of the equiv. relation " \sim "

X/G = partition of X into the orbits of G
($x \sim xg$ or $g(x)$)
for all $g \in G$.

$$X \rightarrow X/\sim$$

$$\mathbb{R}/\mathbb{Z} \cong S^1$$

$$\mathbb{R}^2/\mathbb{Z}^2 \cong T^2 = S^1 \times S^1$$



A non-discrete action of \mathbb{Z} on \mathbb{R} eg. $\{2^k : k \in \mathbb{Z}\}$

