



Math 5605

# Algebraic Topology

Book 3


Cup product for simplicial cohomology  $H^k \times H^l \xrightarrow{\cup} H^{k+l}$   
 makes  $H^*(X; \mathbb{Z})$  or  $H^*(X; \mathbb{R})$  into a graded ring.

To explain, let's talk about singular homology and cohomology.

Singular  $k$ -chains: ( $k = 0, 1, 2, 3, \dots$ ) ways of mapping  $k$ -simplices  
 into  $X$ , not necessarily embeddings.

Take an abstract  $k$ -simplex {all subsets of  $\{0, 1, 2, \dots, k\}$ }.

This has a geometric realization



$$\Delta = \Delta^n = \left\{ \underbrace{(v_0, v_1, \dots, v_n)}_{\text{barycentric coordinates}} : v_i \geq 0, \sum v_i = 1 \right\} \subset \mathbb{R}^{n+1}$$

(convex combinations of  $e_0 = (1, 0, \dots, 0)$ ,  $e_1, \dots, e_n = (0, \dots, 0, 1)$ )

An  $n$ -chain is a formal linear combination of maps  $\sigma: \Delta^n \rightarrow X$ .

$$C_n = \{n\text{-chains in } X\} = C_n(X; \mathbb{R}), \quad \mathbb{R} \text{ any commutative ring with 1} \quad \text{eg. } \mathbb{R}, \mathbb{Z}, \mathbb{F}_2$$

$$C^n = C_n^* = \{n\text{-cochains in } X\} = \text{Hom}(C_n, \mathbb{R}) = \{\mathbb{R}\text{-homomorphisms } C_n \rightarrow \mathbb{R}\}$$

$$d: C_n \rightarrow C_{n-1}, \quad d\sigma = \sum_{i=0}^n \sigma \circ \partial_i \quad d^2 = 0, \quad (d^*)^2 = 0$$

$$d^*: C^{n-1} \rightarrow C^n$$

If  $\phi \in C^k$   $k$ -cochain then  $\phi \cup \psi \in C^{k+l}$  cochain; for any  $(k+l)$ -chain  $\sigma: \Delta^{k+l} \rightarrow X$   
 $\psi \in C^l$   $l$ -cochain  $(\phi \cup \psi)(\sigma) = \phi(\sigma|_{[v_0, \dots, v_k]}) \psi(\sigma|_{[v_k, \dots, v_{k+l}]})$   $[v_0, \dots, v_{k+l}] \mapsto \sigma(v_0, \dots, v_{k+l})$

This gives a bilinear product  $C^k \times C^l \xrightarrow{\cup} C^{k+l}$   
 inducing a bilinear product  $H^k \times H^l \xrightarrow{\cup} H^{k+l}$  (cup product)

making  $H^*(X; \mathbb{R})$  into a graded ring

$$\bigoplus_{i \geq 0} H^i(X; \mathbb{R}).$$

Eg.  $X = \mathbb{P}^n$ ,  $R = \mathbb{F}_2 = \mathbb{Z}/2\mathbb{Z}$  ;  $H^i(X; \mathbb{F}_2) \cong \begin{cases} \mathbb{F}_2 & 0 \leq i \leq n \\ 0 & \text{else} \end{cases}$

$\mathbb{P}^n \mathbb{R} = \{ \text{1-dim'd subspaces of } \mathbb{R}^{n+1} \} = S^n / \text{antipodality}$

$\mathbb{P}^1 \mathbb{R} \cong S^1 / \text{antipodality} \cong S^1$

$\mathbb{O} \cong \mathbb{O}$

$\mathbb{P}^n \mathbb{R}$  is orientable iff  $n$  is odd.

$\mathbb{P}^2 \mathbb{R} = S^2 / \text{antipodality} =$  

$H^*(X; \mathbb{F}_2) \cong \mathbb{F}_2[x] / (x^{n+1})$  Additively:  $\{ a_0 + a_1 x + \dots + a_n x^n : a_i \in \mathbb{F}_2 \}$

Borsuk-Ulam Theorem: There is no antipodal map  $S^n \xrightarrow{f} S^{n-1}$  for  $n \geq 2$ .

Proof is by contradiction


ie.  $f(-x) = -f(x)$

Suppose  $f: S^n \rightarrow S^{n-1}$  is antipodal. ( $f(-x) = -f(x)$ )

Then  $f$  induces a well-defined map

$$\begin{array}{ccc} P^n \mathbb{R} & \xrightarrow{f} & P^{n-1} \mathbb{R} \\ \downarrow \cong & & \downarrow \cong \\ \pm x & & \pm f(x) \end{array}$$

$(x \in S^n)$



$f^*$  maps a generator of  $\pi_1(P^n \mathbb{R})$  to a generator of  $\pi_1(P^{n-1} \mathbb{R})$

$f$  induces  $f^*: H^*(P^n \mathbb{R}; \mathbb{F}_2) \rightarrow H^*(P^{n-1} \mathbb{R}; \mathbb{F}_2)$  mapping  $x \mapsto x$

$$\begin{array}{ccc} \cong & & \cong \\ \mathbb{F}_2[x] / (x^n) & & \mathbb{F}_2[x] / (x^{n-1}) \end{array}$$

$x^n \mapsto x^{n-1}$ ; contradiction.

If  $A$  is an additive abelian gp then  $A \cong \underbrace{\mathbb{Z}^k}_{A/T(A)} \oplus T(A)$  where  $T(A) = \text{torsion subgroup of } A = \{\text{elements of } A \text{ of finite order}\}$

$A/T(A)$  canonically

$k = \text{rank } A = \dim A$ .

For any chain complex  $C_n \xrightarrow{d_n} C_{n-1} \xrightarrow{d_{n-1}} \dots \rightarrow C_2 \xrightarrow{d_2} C_1 \xrightarrow{d_1} C_0 \xrightarrow{0} 0$  (over  $\mathbb{Q}$  or  $\mathbb{R}$ )

we have homology groups  $H_n = \ker d_n / \text{im } d_{n+1}$  with well-defined rank  $H_n(X; \mathbb{Z}) = \text{rank } H_n(X; \mathbb{Q}) = \text{rank } H_n(X; \mathbb{R})$

and Euler characteristic

$$\chi(X) = \sum_{i=0}^n (-1)^i \text{rank } H_i(X) = \sum_{i=0}^n (-1)^i \text{rank } C_i$$

$$C_n \xrightarrow{d_n} C_{n-1} \quad \dim C_n = \dim \ker d_n + \dim \text{im } d_n$$

$$H_n = \ker d_n / \text{im } d_{n+1} \quad \dim H_n = \dim \ker d_n - \dim \text{im } d_{n+1}$$






eg.  $\chi(S^2) = 4 - 6 + 4 = 2$





Closed 2-manifolds i.e. connected compact 2-manifolds without boundary are completely classified using Euler characteristic and orientability (Yes/No)

	$S^2$	$T^2$	$P^2R$	$K^2$
$\dim H_2$	1	1	0	0
$\dim H_1$	0	2	0	1
$\dim H_0$	1	1	1	1
$\chi(X)$	2	0	1	0



$$\chi(S_1 \# S_2) = \chi(S_1) + \chi(S_2) - 2 \quad \text{for any two closed surfaces } S_1, S_2$$

$$\chi(T^2 \# T^2) = \chi(T^2) + \chi(T^2) - 2 = 0 + 0 - 2 = -2$$

$$\underbrace{T^2 \# \dots \# T^2}_g =$$

$$\chi(T^2 \# \dots \# T^2) = 2 - 2g$$

$g =$  genus of orientable surface

$$\chi(P^2R \# P^2R) = 1 + 1 - 2 = 0$$

$K^2$

Exact sequences  $\dots \xrightarrow{d_{n+1}} C_n \xrightarrow{d_n} C_{n-1} \xrightarrow{d_{n-1}} \dots$   $\ker d_n = \text{im } d_{n+1}$

$0 \rightarrow C \rightarrow 0$  is exact  $\iff C = 0$

$0 \rightarrow A \rightarrow B \rightarrow 0$  is exact  $\iff A \cong B$


$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  is exact (short exact)  $\iff C \cong B/A$

If  $f: X \rightarrow X$  is an endomorphism of an abelv. gp.  $X$  (or vector space) (at least in an abelian category) some important short exact sequences are

$$0 \rightarrow \ker f \rightarrow X \xrightarrow{f} f(X) \rightarrow 0$$

$$0 \leftarrow \text{coker } f \leftarrow X \xleftarrow{f} f(X) \leftarrow 0$$

If  $\dots \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  and  $0 \rightarrow C \rightarrow D \rightarrow E \rightarrow \dots$



If  $f: X \rightarrow Y$  then  $\text{coker } f = \varinjlim f(X)$ .  
are exact then we get an exact seq.  
 $\rightarrow A \rightarrow B \rightarrow D \rightarrow E \rightarrow \dots$

$0 \rightarrow \ker f \rightarrow X \xrightarrow{f} X \rightarrow \text{coker } f \rightarrow 0$  is exact.

If  $X$  is a fin. diml vector space over  $F$  then the Euler char. of this sequence is  
 $\dim \text{coker } f - \dim X + \dim X - \dim \ker f = 0$

If  $T: X \rightarrow X$  is an operator (endomorphism) (don't worry about boundedness)  
the index of  $T$  is  $\text{ind } T = \dim \text{coker } T - \dim \ker T$  when both of these terms are finite  
(i.e.  $T$  is Fredholm).

Theorem: Let  $S, T: X \rightarrow X$  be operators (lin. transf).  
 of the three operators  $S, T, ST$ , then whenever two are Fredholm then so is the third and  
 in this case  $\text{ind } ST = \text{ind } S + \text{ind } T$ . (or abd. gps)

In general (i.e. for any lin. transf.  $S, T: X \rightarrow X$ ) we have an exact sequence

$$0 \rightarrow \ker T \rightarrow \ker ST \rightarrow \ker S \rightarrow \text{coker } T \rightarrow \text{coker } ST \rightarrow \text{coker } S \rightarrow 0$$


So its Euler characteristic is zero. i.e.  $\text{ind } S + \text{ind } T - \text{ind } ST = 0$ .

Snake Lemma In an abel. category we have a commutative diagram with exact rows

$$\begin{array}{ccccccc} A & \xrightarrow{f} & B & \xrightarrow{g} & C & \longrightarrow & 0 \\ \downarrow a & & \downarrow b & & \downarrow c & & \\ 0 & \longrightarrow & A' & \xrightarrow{f'} & B' & \xrightarrow{g'} & C' \end{array}$$

then we have a six-term exact seq.

$$\ker a \longrightarrow \ker b \longrightarrow \ker c \longrightarrow \text{coker } a \longrightarrow \text{coker } b \longrightarrow \text{coker } c$$



Group Cohomology: used in the study of group extensions

If  $G$  and  $H$  are groups then an extension of  $H$  by  $G$  is a group  $X$  giving an exact sequence  $1 \rightarrow H \rightarrow X \rightarrow G \rightarrow 1$

Note: Groups are not necessarily abelian. We are asking for a new group  $X$  having a normal subgp  $\cong H$  s.t.  $X/H \cong G$ .  $G$  on top,  $H$  on the bottom.

Trivial:  $X = G \times H$ . (split extension)

$G$  is now an arbitrary group and  $A$  is an abelian group ( $G$  multiplicative;  $A$  additive notation) on which  $G$  acts (each  $g \in G$  gives  $g \in GL(A)$  (automorphisms of  $A$  as an abel. gp or  $\mathbb{Z}$ -module))

$(g_1 g_2)(a) = g_1(g_2(a))$ ;  $g(a+b) = ga + gb$ ;  $1a = a$ .  $G \xrightarrow{\text{homo.}} \text{Aut } A = GL(A)$  (fixed)

We construct an extension of  $A$  by  $G$  i.e. an exact sequence of gps

$$\begin{array}{ccccccc} 1 & \longrightarrow & A & \longrightarrow & \hat{G} & \longrightarrow & G \longrightarrow 1 \\ & & \downarrow & & \downarrow \beta & & \downarrow \gamma \\ 1 & \longrightarrow & A & \longrightarrow & \hat{G} & \longrightarrow & G \longrightarrow 1 \end{array}$$

i.e.  $\hat{G}$  is a gp with normal subgp iso. to  $A$  with  $\hat{G}/A \cong G$

Two extensions  $\hat{G} \hat{\cong} \hat{G}$  are equivalent if we have a commutative diagram as shown with  $\alpha, \beta, \gamma$  isomorphisms of groups (with exact rows), Note that the action of  $G$  on  $A$  is fixed throughout. Cohomology of groups is the tool for this.

...  $\xleftarrow{\delta} C^3 \xleftarrow{\delta} C^2 \xleftarrow{\delta} C^1 \xleftarrow{\delta} C^0 \xleftarrow{\delta} D$  is an exact sequence of additive abel. gps where  $C^k = C^k(G; A)$  is the set of all maps  $G^k \rightarrow A$  as an additive abel. gp. i.e.  $\mathbb{Z}$ -module  
 $\underbrace{G \times G \times \dots \times G}_{k \text{ tuples of } G}$

$$C^0 = A \quad (\text{maps } \{1\} \rightarrow A)$$

$$C^1 = A^G = \text{maps } G \rightarrow A \quad \text{i.e. } f: G \rightarrow A$$

$$C^2 = A^{G \times G} = \text{maps } G \times G \rightarrow A \quad \text{etc.}$$

Given  $a \in C^0$  i.e.  $a \in A$ ,  $\delta a \in C^1$  is  $\delta a: G \rightarrow A$

$$g \mapsto ga - a$$

Given  $f \in C^1$  i.e.  $f: G \rightarrow A$

Construct  $\delta f \in C^2$  i.e.  $(\delta f): G \times G \rightarrow A$

$$(\delta f)(g, h) = \underbrace{g f(h)}_A - \underbrace{f(gh)}_A + \underbrace{f(g)}_A \in A.$$

Given  $f \in C^2$  i.e.  $f: G \times G \rightarrow A$

Construct  $(\delta f): G \times G \times G \rightarrow A$

$$(\delta f)(g, h, k) = g f(h, k) - f(gh, k) + f(g, hk) - f(g, h)$$

See p. 2 bottom of handout for  $\delta: C^k \rightarrow C^{k+1}$  in general

$$\text{Check: } C^2 \xleftarrow{\delta} C^1 \xleftarrow{\delta} C^0 \quad \delta^2 = 0?$$

Take  $a \in C^0 = A$ .

$$(\delta a): G \rightarrow A$$

$$(\delta a)(g) = ga - a.$$

$$\delta^2 a: G \times G \rightarrow A$$

$$(\delta^2 a)(f, g) = f(\delta a)(g) - (\delta a)(fg) + \delta a(f)$$

$$= \cancel{f(ga - a)} - (\cancel{f(g)a} - a) + \cancel{f(a - a)} \\ = \cancel{fga - fa} - \cancel{fga} + a + \cancel{fa - a} \\ = 0$$

Classify extensions  $1 \rightarrow A \rightarrow \hat{G} \rightarrow G \rightarrow 1$  where  $G$  is a group acting on an abelian gp  $A$   
 i.e.  $\hat{G}$  is a group with normal subgp  $A$  with  $\hat{G}/A \cong G$ , using cohomology. Start with a split extension  
 i.e.  $A$  has a complementary subgp in  $\hat{G}$ . So  $\hat{G}$  acts on the subgps complementary to  $A$  by conjugation.  
 $H^1(G; A)$  classifies the complementary subgps up to conjugacy.

Fix an action of  $\underbrace{G}_{\text{mult.}}$  on  $\underbrace{A}_{\text{additively}}$

$$a(g, g_2) = (ag_1)g_2$$

here  $A$  is a right  $G$ -module.

$$a1 = a$$

$\uparrow$  id. of  $G$

$$(a+a')g = ag + a'g$$

for  $a, a' \in A; 1, g, g_1, g_2 \in G$ .

$\hat{G}$  is isomorphic to the

semidirect product  $A \rtimes G = \{ (a, g) : a \in A, g \in G \}$

$$(a_1, g_1)(a_2, g_2) = (a_1g_2 + a_2, g_1g_2)$$

identity  $(0, 1)$

Alternative notation:  $A \rtimes G = \left\{ \begin{bmatrix} g & 0 \\ a & 1 \end{bmatrix} : a \in A, g \in G \right\}$

$\uparrow$  in  $A$

$\uparrow$  in  $G$

$$\begin{bmatrix} g_1 & 0 \\ a_1 & 1 \end{bmatrix} \begin{bmatrix} g_2 & 0 \\ a_2 & 1 \end{bmatrix} = \begin{bmatrix} g_1g_2 & 0 \\ a_1g_2 + a_2 & 1 \end{bmatrix}$$

$$C^2 \xleftarrow{\delta^1} C^1 \xleftarrow{\delta^0} C^0 \xleftarrow{\quad} 0$$

for  $a \in A \stackrel{C^0}{=} C^0$   $(\delta^0)(g) = ag - a$

How do we construct a subgrp of  $A \rtimes G$  complementary to  $A$ ?

Any such subgrp  $H \leq A \rtimes G$  has the form  $\{ (t_g, g) : g \in G \} = \left\{ \begin{bmatrix} g & 0 \\ t_g & 1 \end{bmatrix} : g \in G \right\}$

Here  $g \mapsto t_g, G \rightarrow A$ . This will automatically be a complement to  $A$  as long as it is a subgrp. eg.  $t_1 = 0$  but most importantly, closure.

$$\begin{bmatrix} g & 0 \\ t_g & 1 \end{bmatrix} \begin{bmatrix} g' & 0 \\ t_{g'} & 1 \end{bmatrix} = \begin{bmatrix} gg' & 0 \\ t_{gg'} & 1 \end{bmatrix} \text{ i.e. } \underbrace{t_{gg'}}_A = \underbrace{t_g}_{A} \underbrace{g'}_G + \underbrace{t_{g'}}_A \text{ so } (\delta^1)(g, g') = -f(gg') + f(g)g' + f(g') = 0.$$

Complements of  $A$  in  $\hat{G}$  are given by 1-cocycles.

When are two complements of  $A$  conjugate in  $\hat{G}$ ?  $\hat{G} = A \rtimes G$

If  $A$  has complementary subgps  $H_1, H_2 \leq \hat{G}$  given by  $H_i = \{ (f_i(g), g) : g \in G \}$ ,  $f_i \in Z^1(G; A)$   
 when are  $H_1, H_2$  conjugate in  $\hat{G}$ ?

$$\begin{bmatrix} g & 0 \\ a & 1 \end{bmatrix}^{-1} = \begin{bmatrix} g^{-1} & 0 \\ -ag^{-1} & 1 \end{bmatrix}$$

$$\begin{bmatrix} g & 0 \\ a & 1 \end{bmatrix} \begin{bmatrix} g^{-1} & 0 \\ -ag^{-1} & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Use  $\begin{bmatrix} g & 0 \\ a & 1 \end{bmatrix} \in \hat{G}$  ( $a, g$  fixed) to conjugate  $H_1$ :

$$\begin{bmatrix} g & 0 \\ a & 1 \end{bmatrix} \begin{bmatrix} x & 0 \\ f_1(x) & 1 \end{bmatrix} \begin{bmatrix} g^{-1} & 0 \\ -ag^{-1} & 1 \end{bmatrix} = \begin{bmatrix} gx & 0 \\ ax + f_1(x) & 1 \end{bmatrix} \begin{bmatrix} g^{-1} & 0 \\ -ag^{-1} & 1 \end{bmatrix} = \begin{bmatrix} gxg^{-1} & 0 \\ axg^{-1} + f_1(x)g^{-1} - ag^{-1} & 1 \end{bmatrix} = \begin{bmatrix} gxg^{-1} & 0 \\ f_2(gxg^{-1}) & 1 \end{bmatrix}$$

The 1-cycle defining this conjugate subgroup would have to be  $f_2$ : so

$$f_2(gxg^{-1}) = axg^{-1} + f_1(x)g^{-1} - ag^{-1} = f_2(g)xg^{-1} + f_2(x)g^{-1} - f_2(g)g^{-1}$$

$$\begin{aligned} ax + f_1(x) - a &= f_2(g)x + f_2(x) - f_2(g) \\ f_2(x) - f_1(x) &= (a - f_2(g))x - (a - f_2(g)) \\ &= \delta(a - f_2(g))(x) \end{aligned}$$

$$(\delta f_i)(g, g^{-1}) = f_i(g^{-1}) - f_i(gg^{-1}) + f_i(g)g^{-1} = 0$$

$$\begin{aligned} f_2(gxg^{-1}) &= f_2(gxg^{-1}) \\ &= f_2(g)xg^{-1} + f_2(x)g^{-1} \\ &= f_2(g)xg^{-1} + f_2(x)g^{-1} + f_2(g^{-1}) \\ &= f_2(g)xg^{-1} + f_2(x)g^{-1} - f_2(g)g^{-1} \end{aligned}$$

$f \in C^1$  i.e.  $f: G \rightarrow A$

$$f(xy) = f(x)y + f(y)$$

$$\begin{aligned} f(1) &= f(1 \cdot 1) = f(1) + f(1) \\ &\Rightarrow f(1) = 0 \end{aligned}$$

$$\begin{aligned} 0 &= f(1) = f(gg^{-1}) = f(g)g^{-1} + f(g^{-1}) \\ &\Rightarrow f(g^{-1}) = -f(g)g^{-1} \end{aligned}$$

$f$  is a 1-cocycle:  $f \in Z^1$   
 $(\delta f)(x, y) = f(xy) - f(x)y - f(y) = 0$   
 $f$  is a crossed homomorphism or derivation  
 (If  $G$  acts trivially on  $A$   
 i.e.  $ag = a$  for all  $a \in A, g \in G$ )  
 then  $f$  is a homo.  $G \rightarrow A$ .

Extensions of  $A$  by  $G$  correspond to elements of  $H^1 = Z^1/B^1$ .  
 $f \in B^1$  (1-boundary) iff  
 $f(x) = ax - a = (\delta a)(x)$ ,  $a \in A$ .  
 (principal crossed homomorphisms)  
 (inner derivations)

Ex. Classify extensions of  $C_4 = \langle x : x^4 = 1 \rangle$  by  $C_2 = \langle y : y^2 = 1 \rangle$

$$1 \longrightarrow C_4 \longrightarrow \hat{G} \longrightarrow C_2 \longrightarrow 1$$

Two cases depending on the action of  $C_2$  on  $C_4$

Case I:  $y$  inverts  $x$  i.e.  $yxy^{-1} = x^{-1} = x^3$


$|H'| = 2$ . = how many complementary subgps of  $G$  up to conjugacy.

$C_4$  has four complementary subgps in  $\hat{G} \cong$  dihedral gp of order 8.

$\langle y \rangle, \langle x^2y \rangle$  are conjugate to each other in  $\hat{G}$  ↗ Not conjugate.

$\langle xy \rangle, \langle x^3y \rangle$  are conjugate to each other in  $\hat{G}$  ↖

$$\begin{aligned} x &= (1234) \\ y &= (14)(23) \\ x^2y &= (12)(34) \\ xy &= (13) \\ x^3y &= (24) \end{aligned}$$



Case II:  $y$  commutes with  $x$ .  $xy = yx$   $\hat{G} = C_4 \times C_2$

$\langle x \rangle$  has two complements in  $\hat{G}$ , namely  $\langle y \rangle, \langle x^2y \rangle$ . They are not conjugate.  $|H'| = 2$ .

How many extensions  $1 \longrightarrow C_4 \longrightarrow \hat{G} \longrightarrow C_2 \longrightarrow 1$  are there up to equivalence, if we don't require the extension to be split? (Split  $\iff$  there is a complementary subgp for  $C_4$ )

eg.  $C_8$  is a nonsplit extension of  $C_4$  by  $C_2$ .

Case I:  $C_2$  acts trivially on  $C_4$ . Here there are two extensions:  $C_8$  (nonsplit) and  $C_4 \times C_2$  (split).

Case II:  $C_2$  acts nontrivially on  $C_4$ . Here there are two extensions: dihedral of order 8 (split), quaternion gp of order 8 (nonsplit).



## The Schur-Zassenhaus Theorem

Given groups  $G, N$  with  $G$  acting on  $N$  (the action of  $G$  on  $N$  is fixed) we consider exact sequences of groups

$$1 \longrightarrow N \longrightarrow \hat{G} \longrightarrow G \longrightarrow 1$$

i.e. extensions of  $N$  by  $G$  i.e. groups  $\hat{G}$  having a normal subgroup isomorphic to  $N$ .

If  $|N|, |G|$  are relatively prime then  $H^1 = 1$  and  $H^2 = 1$ .

This says that the extension splits i.e.  $\hat{G}$  has a subgroup complementary to  $N$  and any two complements of  $N$  are conjugate in  $\hat{G}$ .

This generalizes Sylow theory;  $N$  and its complements are Hall subgroups.

Note: We do not require  $N$  to be abelian. If  $N$  is abelian then the formulas are simpler. Even simpler if  $N \subset Z(G)$ .  
(central extension of  $N$  by  $G$ )

A loop is a set  $L$  with a binary operation  $(x, y) \mapsto xy$

such that any two of  $x, y, xy$  uniquely determine the other.

We also assume  $\exists 1 \in L$  such that  $1x = x1 = x$  for all  $x \in L$ .

A Bol loop satisfies  $((xy)z)y = x((yz)y)$  for all  $x, y, z \in L$ .

A Hadamard matrix is an  $n \times n$  matrix  $H$  with entries  $\pm 1$  such that  $HH^T = nI = H^T H$

eg.  $H = \begin{bmatrix} 1 & & & \\ & \oplus & & \\ & & \vdots & \\ & & & \vdots \end{bmatrix}$  gives a (regular) double cover of  $K_{n,n}$ :



A complex Hadamard matrix is an  $n \times n$  matrix  $H$  with entries in  $S = \{z \in \mathbb{C} : |z| = 1\}$  such that  $HH^* = nI = H^* H$

I classified complex Had. matrixes with an automorphism group which is doubly transitive on rows.